

Datasheet: MCA1212F

Description:	RAT IgG2a NEGATIVE CONTROL:FITC
Specificity:	RAT IgG2a NEGATIVE CONTROL
Format:	FITC
Product Type:	Negative/Isotype Control
Isotype:	IgG2a
Quantity:	100 TESTS

Product Details

RRID AB_322675

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/10*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. * It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test antibody.

Target Species Negative Control

Product Form Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Preparation Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative 0.09% Sodium Azide
Stabilisers 1% Bovine Serum Albumin

Approx. Protein Concentrations IgG concentration 0.1 mg/ml

Immunogen Human lymphocytes.

Fusion Partners Spleen cells from immunized DA rats were fused with cells of the rat Y3/Ag1.2.3. myeloma cell line.

Specificity **Rat IgG2a Negative Control antibody** is suitable for the assessment of the level of non-specific binding of rat IgG2a monoclonal antibodies to mouse cells.

Test results indicate Rat IgG2a Negative Control antibody is also suitable for use as a negative control with dog cells.

N.B. This antibody recognizes a human cell surface marker, and therefore is not suitable as a negative control in human cells or cell lines.

References

1. Sumagin, R. *et al.* (2008) Leukocyte-endothelial cell interactions are linked to vascular permeability via ICAM-1-mediated signaling. [Am J Physiol Heart Circ Physiol. 295: H969-H977.](#)
2. Chiu, W.C. *et al.* (2011) Effects of dietary fish oil supplementation on cellular adhesion molecule expression and tissue myeloperoxidase activity in hypercholesterolemic mice with sepsis. [J Nutr Biochem. 20: 254-60.](#)
3. Guilloteau, L.A. *et al.* (2003) Nramp1 is not a major determinant in the control of *Brucella melitensis* infection in mice. [Infect Immun. 71: 621-8.](#)
4. Stapleton, T.W. *et al.* (2000) Investigation of the regenerative capacity of an acellular porcine medial meniscus for tissue engineering applications. [Tissue Eng Part A. 17: 231-42.](#)
5. Park, S.W. *et al.* (2012) A1 adenosine receptor allosteric enhancer PD-81723 protects against renal ischemia-reperfusion injury. [Am J Physiol Renal Physiol. 303: F721-32.](#)
6. Schmidt, E.P. *et al.* (2012) The pulmonary endothelial glycocalyx regulates neutrophil adhesion and lung injury during experimental sepsis. [Nat Med. 18 \(8\): 1217-23.](#)
7. McConnell, M.J. *et al.* (2009) H2-K(b) and H2-D(b) regulate cerebellar long-term depression and limit motor learning. [Proc Natl Acad Sci U S A. 106: 6784-9.](#)
8. Rabadi MM *et al.* (2016) Peptidyl arginine deiminase-4 deficient mice are protected against kidney and liver injury after renal ischemia and reperfusion. [Am J Physiol Renal Physiol. Jun 22: ajrenal.00254.2016. \[Epub ahead of print\]](#)
9. Rabadi, M.M. *et al.* (2019) Peptidyl arginine deiminase-4 exacerbates ischemic AKI by finding NEMO (NFκB Essential Modulator). [Am J Physiol Renal Physiol. Apr 03 \[Epub ahead of print\].](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:FITC \(MCA6005F\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50

'M335591:181204'

Printed on 20 May 2019
