

Datasheet: MCA1212PB

Description:	RAT IgG2a NEGATIVE CONTROL:Pacific Blue®
Specificity:	RAT IgG2a NEGATIVE CONTROL
Format:	Pacific Blue®
Product Type:	Negative/Isotype Control
Isotype:	IgG2a
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures.

Target Species	Negative Control		
Product Form	Purified IgG conjugated to Pacific Blue® - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Pacific Blue®	410	455
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml		
Immunogen	Human lymphocytes.		
RRID	AB_567384		
Fusion Partners	Spleen cells from immunized DA rats were fused with cells of the rat Y3/Ag1.2.3. myeloma cell line.		
Specificity	<p>Rat IgG2a Negative Control antibody is suitable for the assessment of the level of non-specific binding of rat IgG2a monoclonal antibodies to mouse cells.</p> <p>Test results indicate Rat IgG2a Negative Control antibody is also suitable for use as a negative</p>		

control with dog cells.

N.B. This antibody recognizes a human cell surface marker, and therefore is not suitable as a negative control in human cells or cell lines.

Flow Cytometry Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
* It is recommended that the user dilutes the antibody to a concentration equivalent to their test reagent.

References

1. Sumagin, R. *et al.* (2008) Leukocyte-endothelial cell interactions are linked to vascular permeability via ICAM-1-mediated signaling. [Am J Physiol Heart Circ Physiol. 295: H969-H977.](#)
2. Chiu, W.C. *et al.* (2011) Effects of dietary fish oil supplementation on cellular adhesion molecule expression and tissue myeloperoxidase activity in hypercholesterolemic mice with sepsis. [J Nutr Biochem. 20: 254-60.](#)
3. Guilloteau, L.A. *et al.* (2003) Nramp1 is not a major determinant in the control of *Brucella melitensis* infection in mice. [Infect Immun. 71: 621-8.](#)
4. Stapleton, T.W. *et al.* (2000) Investigation of the regenerative capacity of an acellular porcine medial meniscus for tissue engineering applications. [Tissue Eng Part A. 17: 231-42.](#)
5. Park, S.W. *et al.* (2012) A1 adenosine receptor allosteric enhancer PD-81723 protects against renal ischemia-reperfusion injury. [Am J Physiol Renal Physiol. 303: F721-32.](#)
6. Schmidt, E.P. *et al.* (2012) The pulmonary endothelial glycocalyx regulates neutrophil adhesion and lung injury during experimental sepsis. [Nat Med. 18 \(8\): 1217-23.](#)
7. McConnell, M.J. *et al.* (2009) H2-K(b) and H2-D(b) regulate cerebellar long-term depression and limit motor learning. [Proc Natl Acad Sci U S A. 106: 6784-9.](#)
8. Rabadi MM *et al.* (2016) Peptidyl arginine deiminase-4 deficient mice are protected against kidney and liver injury after renal ischemia and reperfusion. [Am J Physiol Renal Physiol. Jun 22: ajprenal.00254.2016. \[Epub ahead of print\]](#)
9. Rabadi, M.M. *et al.* (2019) Peptidyl arginine deiminase-4 exacerbates ischemic AKI by finding NEMO (NFκB Essential Modulator). [Am J Physiol Renal Physiol. Apr 03 \[Epub ahead of print\].](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Acknowledgements The Pacific Blue® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays, and are covered by pending and issued patents.

Health And Safety Information Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:PacificBlue® \(MCA6005PB\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M364935:200529'

Printed on 11 Aug 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)