

Datasheet: MCA1219

Description:	MOUSE ANTI PIG SWC8
Specificity:	SWC8
Format:	S/N
Product Type:	Monoclonal Antibody
Clone:	MIL3
Isotype:	IgM
Quantity:	2 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/20
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Pig
Product Form	Tissue Culture Supernatant - liquid
Preparation	Tissue Culture Supernatant containing 0.2M Tris/HCl pH7.4 and 5-10% foetal calf serum
Preservative Stabilisers	0.09% Sodium Azide
Immunogen	Porcine Lamina Propria Leucocytes.
RRID	AB_322076
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the P3-X63-Ag.653

myeloma cell line.

Specificity **Mouse anti Pig SWC8, clone MIL3**, recognizes the porcine SWC8 cell surface antigen, an antigen that as yet has no identified human homolog. SWC8 is expressed by granulocytes, B cells, a subset of T cells and by some non-haematopoietic cells. Monocytes however do not express SWC8.

Clone MIL3 has been used in two colour flow cytometry with Mouse anti Porcine CD14 antibody, clone MIL2 ([MCA1218GA](#)) to distinguish between monocytes and granulocytes ([Haverson et al. 1994](#)).

Flow Cytometry Use 50ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Haverson, K. *et al.* (1994) Characterization of monoclonal antibodies specific for monocytes, macrophages and granulocytes from porcine peripheral blood and mucosal tissues. [J Immunol Methods. 170 \(2\): 233-45.](#)
2. Lunney, J.K. (1993) Characterization of swine leukocyte differentiation antigens. [Immunol Today. 14 \(4\): 147-8.](#)
3. Summerfield, A. *et al.* (2001) Induction of apoptosis in bone marrow neutrophil-lineage cells by classical swine fever virus. [J Gen Virol. 82 \(Pt 6\): 1309-18.](#)
4. Zelnickova, P. *et al.* (2008) Age-dependent changes of proinflammatory cytokine production by porcine peripheral blood phagocytes. [Vet Immunol Immunopathol. 124: 367-78.](#)
5. Summerfield, A. *et al.* (2003) Porcine peripheral blood dendritic cells and natural interferon-producing cells. [Immunology. 110: 440-9.](#)
6. Barnard, A.L. *et al.* (2005) Immune response characteristics following emergency vaccination of pigs against foot-and-mouth disease. [Vaccine. 23: 1037-47.](#)
7. Chen, L. *et al.* (2003) Macrophages and MHC class II positive dendritiform cells in the iris and choroid of the pig. [Curr Eye Res. 26: 291-6.](#)
8. Ondrackova, P. *et al.* (2010) Porcine mononuclear phagocyte subpopulations in the lung, blood and bone marrow: dynamics during inflammation induced by *Actinobacillus pleuropneumoniae*. [Vet Res. 41: 64.](#)
9. LeLuduec, J.B. *et al.* (2016) Intradermal vaccination with un-adjuvanted sub-unit vaccines triggers skin innate immunity and confers protective respiratory immunity in domestic swine. [Vaccine. 34 \(7\): 914-22.](#)

Further Reading 1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10053 available at:
10053: <https://www.bio-rad-antibodies.com/uploads/MSDS/10053.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgM (STAR138...) [Alk. Phos.](#)
Human Anti Mouse IgM (HCA040...) [FITC](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgM NEGATIVE CONTROL \(MCA692\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M383205:210513'

Printed on 29 Aug 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)