

Datasheet: MCA1222F

Description:	MOUSE ANTI PIG CD45:FITC
Specificity:	CD45
Other names:	LCA
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	K252.1E4
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% sodium azide (NaN ₃)		
Stabilisers	1% bovine serum albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		

Immunogen	Porcine Peripheral Blood lymphocytes.
RRID	AB_323237
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the P3 - X63 - Ag.653 myeloma cell line.
Specificity	<p>Mouse anti Pig CD45, clone K252.1E4 recognizes an epitope common to all porcine CD45 isoforms (Schnitzlein et al. 1998). CD45 is also known as leukocyte common antigen (LCA).</p> <p>Mouse anti Pig CD45, clone K252.1E4 immunoprecipitates three polypeptides of 226, 210 and 190 kDa from preparations of porcine peripheral blood mononuclear cells and shows a broad reactivity pattern with both lymphoid and myeloid cells (Zuckermann et al. 1994).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> 1. Yang, P. <i>et al.</i> (2002) Immune cells in the porcine retina: distribution, characterization and morphological features. Invest Ophthalmol Vis Sci. 43 (5): 1488-92. 2. Terzic, S. <i>et al.</i> (2002) Immunophenotyping of leukocyte subsets in peripheral blood and palatine tonsils of prefattening pigs. Vet Res Commun. 26: 273-83. 3. Barker, E. <i>et al.</i> (2006) The larynx as an immunological organ: immunological architecture in the pig as a large animal model. Clin Exp Immunol. 143: 6-14. 4. Vilahur, G. <i>et al.</i> (2015) Roflumilast-induced Local Vascular Injury Is Associated with a Coordinated Proteome and Microparticle Change in the Systemic Circulation in Pigs. Toxicol Pathol. 43 (4): 569-80. 5. O'Leary, S. <i>et al.</i> (2004) Seminal plasma regulates endometrial cytokine expression, leukocyte recruitment and embryo development in the pig. Reproduction. 128: 237-47. 6. Zelnickova, P. <i>et al.</i> (2006) Postnatal functional maturation of blood phagocytes in pig. Vet Immunol Immunopathol. 113: 383-91. 7. Bimczok, D. <i>et al.</i> (2006) Phenotype and distribution of dendritic cells in the porcine small intestinal and tracheal mucosa and their spatial relationship to epithelial cells. Cell Tissue Res. 325: 461-8. 8. Nochi, T. <i>et al.</i> (2004) Biological role of Ep-CAM in the physical interaction between epithelial cells and lymphocytes in intestinal epithelium. Clin Immunol. 113: 326-39. 9. Bimczok, D. <i>et al.</i> (2010) Primary porcine CD11R1+ antigen-presenting cells isolated from small intestinal mucosa mature but lose their T cell stimulatory function in response to cholera toxin treatment. Vet Immunol Immunopathol. 134: 239-48. 10. Ebdrup, L. <i>et al.</i> (2008) Dynamic expression of the signal regulatory protein alpha and CD18 on porcine PBMC during acute endotoxaemia. Scand J Immunol. 68: 430-7. 11. Plánka, L. <i>et al.</i> (2009) Use of allogenic stem cells for the prevention of bone bridge formation in miniature pigs. Physiol Res. 58: 885-93. 12. Plánka, L. <i>et al.</i> (2009) Comparison of Preventive and Therapeutic Transplantations of Allogeneic Mesenchymal Stem Cells in Healing of the Distal Femoral Growth Plate Cartilage Defects in Miniature Pigs. Acta Vet. Brno 78: 293-302. 13. Plánka, L. <i>et al.</i> (2008) New options for management of posttraumatic articular cartilage defects. Rozhl Chir. 87: 42-5. 14. Splíchal, I. <i>et al.</i> (2002) <i>Escherichia coli</i> Administered into Pig Amniotic Cavity Appear

- in Fetal Airways and Attract Macrophages into Fetal Lungs. [Physiol Res. 51: 523-8.](#)
15. Juhászová, J. *et al.* (2011) Osteogenic differentiation of miniature pig mesenchymal stem cells in 2D and 3D environment. [Physiol Res. 60: 559-71.](#)
16. Heino, T.J. *et al.* (2012) Comparison of the osteogenic capacity of minipig and human bone marrow-derived mesenchymal stem cells. [J Orthop Res. 30: 1019-25.](#)
17. Stepanova, H. *et al.* (2012) CD4⁺ and $\gamma\delta$ TCR⁺ T lymphocytes are sources of interleukin-17 in swine. [Cytokine. 58: 152-7.](#)
18. Hester, S.N. *et al.* (2012) Intestinal and systemic immune development and response to vaccination are unaffected by dietary (1,3/1,6)- β -D-glucan supplementation in neonatal piglets. [Clin Vaccine Immunol. 19: 1499-508.](#)
19. Noort, W.A. *et al.* (2011) Human versus porcine mesenchymal stromal cells: phenotype, differentiation potential, immunomodulation and cardiac improvement after transplantation. [J Cell Mol Med. 16: 1827-39.](#)
20. Thorum, S.C. *et al.* (2013) Impact of diet on development of bronchial-associated immunity in the neonatal piglet. [Vet Immunol Immunopathol. 151: 63-72.](#)
21. Tran, L.A. *et al.* (2010) Gadonanotubes as Magnetic Nanolabels for Stem Cell Detection [Biomaterials. 31: 9482-91.](#)
22. Thorum, S.C. *et al.* (2013) Dietary (1,3/1,6)- β -D-glucan decreases transforming growth factor β expression in the lung of the neonatal piglet. [Nutr Res. 33 \(4\): 322-31.](#)
23. Koskinas, K.C. *et al.* (2013) Synergistic effect of local endothelial shear stress and systemic hypercholesterolemia on coronary atherosclerotic plaque progression and composition in pigs. [Int J Cardiol. 169 \(6\): 394-401.](#)
24. Franzoni, G. *et al.* (2013) Assessment of the Phenotype and Functionality of Porcine CD8 T Cell Responses following Vaccination with Live Attenuated Classical Swine Fever Virus (CSFV) and Virulent CSFV Challenge. [Clin Vaccine Immunol. 20: 1604-16.](#)
25. Filová E, *et al.* (2009) Regionally-selective cell colonization of micropatterned surfaces prepared by plasma polymerization of acrylic acid and 1,7-octadiene. [Physiol Res. 58: 669-84.](#)
26. Bimczok, D. *et al.* (2005) Site-specific expression of CD11b and SIRPalpha (CD172a) on dendritic cells: implications for their migration patterns in the gut immune system. [Eur J Immunol. 35: 1418-27.](#)
27. Sun, Z. *et al.* (2013) Scaffold-Based Delivery of Autologous Mesenchymal Stem Cells for Mandibular Distraction Osteogenesis: Preliminary Studies in a Porcine Model. [PLoS One. 8: e74672.](#)
28. Post, I.C. *et al.* (2013) Characterization and quantification of porcine circulating endothelial cells. [Xenotransplantation. 20: 18-26.](#)
29. Foubert, P. *et al.* (2015) Uncultured adipose-derived regenerative cells (ADRCs) seeded in collagen scaffold improves dermal regeneration, enhancing early vascularization and structural organization following thermal burns. [Burns. 41 \(7\): 1504-16.](#)
30. Rayat, G.R. *et al.* (2016) First update of the International Xenotransplantation Association consensus statement on conditions for undertaking clinical trials of porcine islet products in type 1 diabetes--Chapter 3: Porcine islet product manufacturing and release testing criteria. [Xenotransplantation. 23 \(1\): 38-45.](#)
31. Antonson, A.M. *et al.* (2017) Maternal viral infection during pregnancy elicits anti-social behavior in neonatal piglet offspring independent of postnatal microglial cell activation. [Brain Behav Immun. 59: 300-312.](#)

32. Westover, A.J. *et al.* (2016) An Immunomodulatory Device Improves Insulin Resistance in Obese Porcine Model of Metabolic Syndrome. [J Diabetes Res. 2016: 3486727.](#)
33. Kaczmarek, M.M. *et al.* (2013) Seminal plasma affects prostaglandin synthesis and angiogenesis in the porcine uterus. [Biol Reprod. 88 \(3\): 72.](#)
34. Valpotić, H. *et al.* (2018) Dietary supplementation with mannan oligosaccharide and clinoptilolite modulates innate and adaptive immune parameters of weaned pigs. [Pol J Vet Sci. 21 \(1\): 83-93.](#)
35. Sauerova, P. *et al.* (2019) Positive impact of dynamic seeding of mesenchymal stem cells on bone-like biodegradable scaffolds with increased content of calcium phosphate nanoparticles. [Mol Biol Rep. 46 \(4\): 4483-4500.](#)
36. McCleary, S. *et al.* (2020) Substitution of warthog NF- κ B motifs into RELA of domestic pigs is not sufficient to confer resilience to African swine fever virus. [Sci Rep. 10 \(1\): 8951.](#)
37. Zhang, W. *et al.* (2020) Dietary Calcium and Phosphorus Amounts Affect Development and Tissue-Specific Stem Cell Characteristics in Neonatal Pigs. [J Nutr. 150 \(5\): 1086-92.](#)
38. Teuben, M.P.J. *et al.* (2021) Standardized porcine unilateral femoral nailing is associated with changes in PMN activation status, rather than aberrant systemic PMN prevalence. [Eur J Trauma Emerg Surg. Jun 10 \[Epub ahead of print\].](#)
39. McCormack, U.M. *et al.* (2019) Porcine Feed Efficiency-Associated Intestinal Microbiota and Physiological Traits: Finding Consistent Cross-Locational Biomarkers for Residual Feed Intake. [mSystems.4 \(4\): e00324-18.](#)
40. Zimmermann, C.E. *et al.* (2021) Characterization of porcine mesenchymal stromal cells and their proliferative and osteogenic potential in long-term culture. [J Stem Cells Regen Med. 17 \(2\): 49-55.](#)
41. Zhu, H. *et al.* (2022) Production of cultured meat from pig muscle stem cells. [Biomaterials. 287: 121650.](#)

Further Reading 1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M408573:221013'

Printed on 13 Oct 2022

© 2022 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)