

Datasheet: MCA1226FT

BATCH NUMBER 173724

Description:	MOUSE ANTI HUMAN CD8:FITC
Specificity:	CD8
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	LT8
Isotype:	IgG1
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/5

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Marmoset, Chimpanzee, Cynomolgus monkey, Red-bellied Tamarin
N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

Preparation

Purified IgG prepared by ion exchange chromatography from ascites

Buffer Solution

Phosphate buffered saline

Preservative	0.09% sodium azide (NaN ₃)
Stabilisers	1% bovine serum albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Normal human peripheral blood lymphocytes.
External Database Links	<p>UniProt:</p> <p>P01732 Related reagents</p> <p>P10966 Related reagents</p> <p>Entrez Gene:</p> <p>925 CD8A Related reagents</p> <p>926 CD8B Related reagents</p>
Synonyms	CD8B1, MAL
RRID	AB_1102345
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63.653 myeloma cell line.
Specificity	<p>Mouse anti Human CD8 antibody, clone LT8 recognizes the human CD8 cell surface glycoprotein expressed by a subset of peripheral blood T cells which express cytotoxic/suppressor activity. It is also expressed weakly on NK cells.</p> <p>The CD8 antigen is a co-receptor for MHC Class I in conjunction with the T cell receptor, and is important in the selection process of CD8+ MHC Class I restricted T cells.</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells or cells or 100µl human whole blood
References	<ol style="list-style-type: none"> Zarkesh-Esfahani, H. <i>et al.</i> (2001) High-dose leptin activates human leukocytes via receptor expression on monocytes. J Immunol. 167 (8): 4593-9. Manninen, A. & Saksela, K. (2002) HIV-1 Nef interacts with inositol trisphosphate receptor to activate calcium signaling in T cells. J Exp Med. 195 (8): 1023-32. Parnes, J.R. (1989) Molecular biology and function of CD4 and CD8. Adv Immunol. 44: 265-311. Kap, Y.S. <i>et al.</i> (2009) A monoclonal antibody selection for immunohistochemical examination of lymphoid tissues from non-human primates. J Histochem Cytochem. 57: 1159-67. Hovden, A.O. <i>et al.</i> (2011) Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. BMC Immunol. 12: 2. Nelson, M. <i>et al.</i> (2010) Characterization of lethal inhalational infection with <i>Francisella tularensis</i> in the common marmoset (<i>Callithrix jacchus</i>). J Med Microbiol. 59: 1107-13. Gibbings, D.J. <i>et al.</i> (2007) CD8 alpha is expressed by human monocytes and

- enhances Fc gamma R-dependent responses. [BMC Immunol. 8: 12.](#)
8. Junker, A. *et al.* (2007) Multiple sclerosis: T-cell receptor expression in distinct brain regions. [Brain. 130: 2789-99.](#)
9. Held, K. *et al.* (2011) Expression of herpes simplex virus 1-encoded microRNAs in human trigeminal ganglia and their relation to local T-cell infiltrates. [J Virol. 85 \(19\): 9680-5.](#)
10. Hood SP *et al.* (2014) Changes in immune cell populations in the periphery and liver of GBV-B-infected and convalescent tamarins (*Saguinus labiatus*). [Virus Res. 179: 93-101.](#)
11. Nelson, M. & Loveday, M. (2014) Exploring the innate immunological response of an alternative nonhuman primate model of infectious disease; the common marmoset. [J Immunol Res. 2014: 913632.](#)
12. Manivannan, K. *et al.* (2016) CADM1/TSLC1 Identifies HTLV-1-Infected Cells and Determines Their Susceptibility to CTL-Mediated Lysis. [PLoS Pathog. 12 \(4\): e1005560.](#)
13. Gross, C.C. *et al.* (2016) Impaired NK-mediated regulation of T-cell activity in multiple sclerosis is reconstituted by IL-2 receptor modulation. [Proc Natl Acad Sci U S A. 113 \(21\): E2973-82.](#)
14. Dunham, J. *et al.* (2016) Blockade of CD127 Exerts a Dichotomous Clinical Effect in Marmoset Experimental Autoimmune Encephalomyelitis. [J Neuroimmune Pharmacol. 11 \(1\): 73-83.](#)
15. Bughani, U. *et al.* (2017) T cell activation and differentiation is modulated by a CD6 domain 1 antibody Itolizumab. [PLoS One. 12 \(7\): e0180088.](#)
16. Philippens, I.H. *et al.* (2017) Acceleration of Amyloidosis by Inflammation in the Amyloid-Beta Marmoset Monkey Model of Alzheimer's Disease. [J Alzheimers Dis. 55 \(1\): 101-113.](#)
17. Flak, V.R. *et al.* (2024) Pembrolizumab followed by irreversible electroporation of a liver metastasis in pancreatic cancer patients [iScience. 27 \(10\): 111026.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1226FT>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M437015:250307'

Printed on 28 May 2026

© 2026 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)