

## Datasheet: MCA1226SBB700

**BATCH NUMBER 100006327**

<b>Description:</b>	MOUSE ANTI HUMAN CD8:StarBright Blue 700
<b>Specificity:</b>	CD8
<b>Format:</b>	StarBright Blue 700
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	LT8
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS/0.5ml

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

#### Target Species

Human

#### Species Cross Reactivity

Reacts with: Marmoset, Chimpanzee, Cynomolgus monkey, Red-bellied Tamarin  
**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

#### Product Form

Purified IgG conjugated to StarBright Blue 700 - liquid

#### Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
StarBright Blue 700	470	700

#### Preparation

Purified IgG prepared by ion exchange chromatography

#### Buffer Solution

Phosphate buffered saline

<b>Preservative</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Stabilisers</b>	1% Bovine Serum Albumin 0.1% Pluronic F68 0.1% PEG 3350 0.05% Tween 20
<b>Immunogen</b>	Normal human peripheral blood lymphocytes.
<b>External Database Links</b>	<p><b>UniProt:</b></p> <p><a href="#">P01732</a>    <a href="#">Related reagents</a></p> <p><a href="#">P10966</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b></p> <p><a href="#">925</a> CD8A    <a href="#">Related reagents</a></p> <p><a href="#">926</a> CD8B    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	CD8B1, MAL
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63.653 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Human CD8 antibody, clone LT8</b> recognizes the human CD8 cell surface glycoprotein expressed by a subset of peripheral blood T cells which express cytotoxic/suppressor activity. It is also expressed weakly on NK cells.</p> <p>The CD8 antigen is a co-receptor for MHC Class I in conjunction with the T cell receptor, and is important in the selection process of CD8+ MHC Class I restricted T cells.</p>
<b>Flow Cytometry</b>	Use 5ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
<b>References</b>	<ol style="list-style-type: none"> <li>Zarkesh-Esfahani, H. <i>et al.</i> (2001) High-dose leptin activates human leukocytes via receptor expression on monocytes. <a href="#">J Immunol. 167 (8): 4593-9.</a></li> <li>Manninen, A. &amp; Saksela, K. (2002) HIV-1 Nef interacts with inositol trisphosphate receptor to activate calcium signaling in T cells. <a href="#">J Exp Med. 195 (8): 1023-32.</a></li> <li>Parnes, J.R. (1989) Molecular biology and function of CD4 and CD8. <a href="#">Adv Immunol. 44: 265-311.</a></li> <li>Kap, Y.S. <i>et al.</i> (2009) A monoclonal antibody selection for immunohistochemical examination of lymphoid tissues from non-human primates. <a href="#">J Histochem Cytochem. 57: 1159-67.</a></li> <li>Hovden, A.O. <i>et al.</i> (2011) Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. <a href="#">BMC Immunol. 12: 2.</a></li> <li>Nelson, M. <i>et al.</i> (2010) Characterization of lethal inhalational infection with <i>Francisella tularensis</i> in the common marmoset (<i>Callithrix jacchus</i>). <a href="#">J Med Microbiol. 59: 1107-13.</a></li> <li>Gibbings, D.J. <i>et al.</i> (2007) CD8 alpha is expressed by human monocytes and enhances Fc gamma R-dependent responses. <a href="#">BMC Immunol. 8: 12.</a></li> <li>Junker, A. <i>et al.</i> (2007) Multiple sclerosis: T-cell receptor expression in distinct brain</li> </ol>

regions. [Brain. 130: 2789-99.](#)

9. Held, K. *et al.* (2011) Expression of herpes simplex virus 1-encoded microRNAs in human trigeminal ganglia and their relation to local T-cell infiltrates. [J Virol. 85 \(19\): 9680-5.](#)

10. Hood SP *et al.* (2014) Changes in immune cell populations in the periphery and liver of GBV-B-infected and convalescent tamarins (*Saguinus labiatus*). [Virus Res. 179: 93-101.](#)

11. Nelson, M. & Loveday, M. (2014) Exploring the innate immunological response of an alternative nonhuman primate model of infectious disease; the common marmoset. [J Immunol Res. 2014: 913632.](#)

12. Manivannan, K. *et al.* (2016) CADM1/TSLC1 Identifies HTLV-1-Infected Cells and Determines Their Susceptibility to CTL-Mediated Lysis. [PLoS Pathog. 12 \(4\): e1005560.](#)

13. Gross, C.C. *et al.* (2016) Impaired NK-mediated regulation of T-cell activity in multiple sclerosis is reconstituted by IL-2 receptor modulation. [Proc Natl Acad Sci U S A. 113 \(21\): E2973-82.](#)

14. Dunham, J. *et al.* (2016) Blockade of CD127 Exerts a Dichotomous Clinical Effect in Marmoset Experimental Autoimmune Encephalomyelitis. [J Neuroimmune Pharmacol. 11 \(1\): 73-83.](#)

15. Bughani, U. *et al.* (2017) T cell activation and differentiation is modulated by a CD6 domain 1 antibody Itolizumab. [PLoS One. 12 \(7\): e0180088.](#)

16. Philippens, I.H. *et al.* (2017) Acceleration of Amyloidosis by Inflammation in the Amyloid-Beta Marmoset Monkey Model of Alzheimer's Disease. [J Alzheimers Dis. 55 \(1\): 101-113.](#)

---

<b>Storage</b>	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
<b>Guarantee</b>	12 months from date of despatch
<b>Acknowledgements</b>	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #20471 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1226SBB700">https://www.bio-rad-antibodies.com/SDS/MCA1226SBB700</a> 20471
<b>Regulatory</b>	For research purposes only

---

## Related Products

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South America** Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](http://bio-rad-antibodies.com/datasheets)

