

Datasheet: MCA1226SBUV510

Description:	MOUSE ANTI HUMAN CD8:StarBright UltraViolet 510		
Specificity:	CD8		
Format:	StarBright UltraViolet 510		
Product Type:	Monoclonal Antibody		
Clone:	LT8		
Isotype:	lgG1		
Quantity:	100 TESTS/0.5ml		

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .				
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry	-			Neat
	Where this product ha	as not been teste	ed for u	se in a particular tech	nique this does not
	necessarily exclude it a guide only. It is reco system using appropr	ommended that t	the user	titrates the product for	g dilutions are given as or use in their own
Target Species	Human				
Species Cross Reactivity	Reacts with: Marmose N.B. Antibody reactivi reactivity is derived fro personal communicati further information.	ity and working o om testing withir	condition n our lat	ns may vary between poratories, peer-reviev	species. Cross wed publications or
Product Form	Purified IgG conjugated to StarBright UltraViolet 510 - liquid				
Max Ex/Em	Fluorophore	Excitation Max	(nm)	Emission Max (nm)	
	StarBright UltraViolet 510	340		513	
Preparation	Purified IgG prepared	by ion exchang	e chron	natography from ascit	es
Buffer Solution	Phosphate buffered sa	aline			
Preservative	0.09% sodium azide ((NaN ₃)			

Stabilisers	1% bovine serum albumin 0.1% Pluronic F68 0.1% PEG 3350 0.05% Tween 20
Immunogen	Normal human peripheral blood lymphocytes.
External Database Links	UniProt:
Links	P01732 Related reagents
	P10966 Related reagents
	Entros Conos
	Entrez Gene: 925 CD8A <u>Related reagents</u>
	926 CD8B Related reagents
Synonyms	CD8B1, MAL
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63.653 myeloma cell line.
Specificity	Mouse anti Human CD8 antibody, clone LT8 recognizes the human CD8 cell surface glycoprotein expressed by a subset of peripheral blood T cells which express cytotoxic/suppressor activity. It is also expressed weakly on NK cells.
	The CD8 antigen is a co-receptor for MHC Class I in conjunction with the T cell receptor, and is important in the selection process of CD8+ MHC Class I restricted T cells.
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 ⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
References	1. Zarkesh-Esfahani, H. <i>et al.</i> (2001) High-dose leptin activates human leukocytes via receptor expression on monocytes. <u>J Immunol. 167 (8): 4593-9.</u>
	2. Manninen, A. & Saksela, K. (2002) HIV-1 Nef interacts with inositol trisphosphate
	receptor to activate calcium signaling in T cells. <u>J Exp Med. 195 (8): 1023-32.</u>
	3. Parnes, J.R. (1989) Molecular biology and function of CD4 and CD8. <u>Adv Immunol. 44:</u> <u>265-311.</u>
	4. Kap, Y.S. <i>et al.</i> (2009) A monoclonal antibody selection for immunohistochemical
	examination of lymphoid tissues from non-human primates. <u>J Histochem Cytochem. 57:</u>
	<u>1159-67.</u> 5. Hovden, A.O. <i>et al.</i> (2011) Maturation of monocyte derived dendritic cells with OK432
	boosts IL-12p70 secretion and conveys strong T-cell responses. <u>BMC Immunol. 12: 2.</u>
	6. Nelson, M. et al. (2010) Characterization of lethal inhalational infection with Francisella
	<i>tularensis</i> in the common marmoset (<i>Callithrix jacchus</i>). <u>J Med Microbiol. 59: 1107-13.</u>
	7. Gibbings, D.J. <i>et al.</i> (2007) CD8 alpha is expressed by human monocytes and
	enhances Fc gamma R-dependent responses. <u>BMC Immunol. 8: 12.</u>
	8. Junker, A. <i>et al.</i> (2007) Multiple sclerosis: T-cell receptor expression in distinct brain regions. <u>Brain. 130: 2789-99.</u>
	·

	 9. Held, K. <i>et al.</i> (2011) Expression of herpes simplex virus 1-encoded microRNAs in human trigeminal ganglia and their relation to local T-cell infiltrates. J Virol. 85 (19): 9680-5. 10. Hood SP <i>et al.</i> (2014) Changes in immune cell populations in the periphery and liver of GBV-B-infected and convalescent tamarins (<i>Saguinus labiatus</i>). Virus Res. 179: 93-101. 11. Nelson, M. & Loveday, M. (2014) Exploring the innate immunological response of an alternative nonhuman primate model of infectious disease; the common marmoset. J Immunol Res. 2014: 913632. 12. Manivannan, K. <i>et al.</i> (2016) CADM1/TSLC1 Identifies HTLV-1-Infected Cells and Determines Their Susceptibility to CTL-Mediated Lysis. PLoS Pathog. 12 (4): e1005560. 13. Gross, C.C. <i>et al.</i> (2016) Impaired NK-mediated regulation of T-cell activity in multiple sclerosis is reconstituted by IL-2 receptor modulation. Proc Natl Acad Sci U S A. 113 (21): E2973-82. 14. Dunham, J. <i>et al.</i> (2016) Blockade of CD127 Exerts a Dichotomous Clinical Effect in Marmoset Experimental Autoimmune Encephalomyelitis. J Neuroimmune Pharmacol. 11 (1): 73-83. 15. Bughani, U. <i>et al.</i> (2017) T cell activation and differentiation is modulated by a CD6 domain 1 antibody Itolizumab. PLoS One. 12 (7): e0180088. 16. Philippens, I.H. <i>et al.</i> (2017) Acceleration of Amyloidosis by Inflammation in the Amyloid-Beta Marmoset Monkey Model of Alzheimer's Disease. J Alzheimers Dis. 55 (1): 101-113.
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts

Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA1226SBUV510 20471
Regulatory	For research purposes only

For research purposes only

Related Products

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide
America	Fax: +1 919 878 3751	
	Email: antibody_sales_us@bio-rad.c	com

Tel: +44 (0)1865 852 700 Europe Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint