

Datasheet: MCA1226SBUV575

Description:	MOUSE ANTI HUMAN CD8:StarBright UltraViolet 575
Specificity:	CD8
Format:	StarBright UltraViolet 575
Product Type:	Monoclonal Antibody
Clone:	LT8
Isotype:	IgG1
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Reacts with: Marmoset, Chimpanzee, Cynomolgus monkey, Rec N.B. Antibody reactivity and working conditions may vary between

Reacts with: Marmoset, Chimpanzee, Cynomolgus monkey, Red-bellied Tamarin **N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form	Purified IgG conjugated to StarBright UltraViolet 575 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	StarBright UltraViolet 575	340	569
Preparation	Purified IgG prepared	d by ion exchange chro	matography from as

- ,	r armou igo proparou sy ion oxonango omomatograpny nom at
Buffer Solution	Phosphate buffered saline
Preservative	0.09% sodium azide (NaN ₃)

Stabilisers

1% bovine serum albumin

0.1% Pluronic F68 0.1% PEG 3350 0.05% Tween 20

Immunogen

Normal human peripheral blood lymphocytes.

External Database Links

UniProt:

P01732 Related reagents
P10966 Related reagents

Entrez Gene:

925CD8ARelated reagents926CD8BRelated reagents

Synonyms

CD8B1, MAL

Fusion Partners

Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63.653 myeloma cell line.

Specificity

Mouse anti Human CD8 antibody, clone LT8 recognizes the human CD8 cell surface glycoprotein expressed by a subset of peripheral blood T cells which express cytotoxic/suppressor activity. It is also expressed weakly on NK cells.

The CD8 antigen is a co-receptor for MHC Class I in conjunction with the T cell receptor, and is important in the selection process of CD8+ MHC Class I restricted T cells.

Flow Cytometry

Use 5μ I of the suggested working dilution to label 10^6 cells in 100μ I. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

References

- 1. Zarkesh-Esfahani, H. *et al.* (2001) High-dose leptin activates human leukocytes via receptor expression on monocytes. <u>J Immunol. 167 (8): 4593-9.</u>
- 2. Manninen, A. & Saksela, K. (2002) HIV-1 Nef interacts with inositol trisphosphate receptor to activate calcium signaling in T cells. J Exp Med. 195 (8): 1023-32.
- 3. Parnes, J.R. (1989) Molecular biology and function of CD4 and CD8. <u>Adv Immunol. 44:</u> 265-311.
- 4. Kap, Y.S. *et al.* (2009) A monoclonal antibody selection for immunohistochemical examination of lymphoid tissues from non-human primates. <u>J Histochem Cytochem. 57:</u> 1159-67.
- 5. Hovden, A.O. *et al.* (2011) Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. <u>BMC Immunol.</u> 12: 2.
- 6. Nelson, M. *et al.* (2010) Characterization of lethal inhalational infection with *Francisella tularensis* in the common marmoset (*Callithrix jacchus*). <u>J Med Microbiol. 59: 1107-13.</u>
- 7. Gibbings, D.J. *et al.* (2007) CD8 alpha is expressed by human monocytes and enhances Fc gamma R-dependent responses. BMC Immunol. 8: 12.
- 8. Junker, A. *et al.* (2007) Multiple sclerosis: T-cell receptor expression in distinct brain regions. Brain. 130: 2789-99.

- 9. Held, K. *et al.* (2011) Expression of herpes simplex virus 1-encoded microRNAs in human trigeminal ganglia and their relation to local T-cell infiltrates. <u>J Virol. 85 (19):</u> 9680-5.
- 10. Hood SP *et al.* (2014) Changes in immune cell populations in the periphery and liver of GBV-B-infected and convalescent tamarins (*Saguinus labiatus*). <u>Virus Res. 179:</u> 93-101.
- 11. Nelson, M. & Loveday, M. (2014) Exploring the innate immunological response of an alternative nonhuman primate model of infectious disease; the common marmoset. <u>J. Immunol Res. 2014: 913632.</u>
- 12. Manivannan, K. *et al.* (2016) CADM1/TSLC1 Identifies HTLV-1-Infected Cells and Determines Their Susceptibility to CTL-Mediated Lysis. PLoS Pathog. 12 (4): e1005560.
- 13. Gross, C.C. *et al.* (2016) Impaired NK-mediated regulation of T-cell activity in multiple sclerosis is reconstituted by IL-2 receptor modulation. <u>Proc Natl Acad Sci U S A. 113 (21):</u> E2973-82.
- 14. Dunham, J. *et al.* (2016) Blockade of CD127 Exerts a Dichotomous Clinical Effect in Marmoset Experimental Autoimmune Encephalomyelitis. <u>J Neuroimmune Pharmacol. 11</u> (1): 73-83.
- 15. Bughani, U. *et al.* (2017) T cell activation and differentiation is modulated by a CD6 domain 1 antibody Itolizumab. <u>PLoS One. 12 (7): e0180088.</u>
- 16. Philippens, I.H. *et al.* (2017) Acceleration of Amyloidosis by Inflammation in the Amyloid-Beta Marmoset Monkey Model of Alzheimer's Disease. <u>J Alzheimers Dis. 55 (1):</u> 101-113.

Storage	Store at +4°C. DO NOT FREEZE.
	This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and fore counterparts
Health And Safety	Material Safety Datasheet documentation #20471 available at:
Information	https://www.bio-rad-antibodies.com/SDS/MCA1226SBUV575
	20471
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M410657:221028'

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint