

Datasheet: MCA1226SBUV795

### **BATCH NUMBER 100006204**

| Description:  | MOUSE ANTI HUMAN CD8:StarBright UltraViolet 795 |
|---------------|---|
| Specificity:  | CD8   |
| Format:       | StarBright UltraViolet 795                      |
| Product Type: | Monoclonal Antibody                             |
| Clone:        | LT8   |
| Isotype:      | lgG1  |
| Quantity:     | 100 TESTS/0.5ml                                 |
|               |   |

# **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry |     |    |                | Neat               |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| Target Species              | Human  |   |  |  |
|-----------------------------|--|---|--|--|
| Species Cross<br>Reactivity | <b>N.B.</b> Antibody reactive reactivity is derived from | et, Chimpanzee, Cyno<br>rity and working conditi<br>rom testing within our la<br>tions from the originato | ons may vary between aboratories, peer-rev | en species. Cross<br>riewed publications o |
| Product Form                | Purified IgG conjugat                                    | ed to StarBright UltraV   | iolet 795 - liquid                         |  |
| Max Ex/Em                   | Fluorophore  | Excitation Max (nm)   | Emission Max (nm)                          |  |
|                             | StarBright UltraViolet<br>795                            | 340   | 792  |  |
| Preparation                 | Purified IgG prepared                                    | d by ion exchange chro  | omatography                                |  |
| Buffer Solution             | Phosphate buffered s                                     | saline  |  |  |

| P | re | se  | rv  | at | ive | • |
|---|----|-----|-----|----|-----|---|
| S | ta | bil | lis | er | s   |   |

0.09% Sodium Azide (NaN<sub>3</sub>)1% Bovine Serum Albumin

0.1% Pluronic F68 0.1% PEG 3350 0.05% Tween 20

#### Immunogen

Normal human peripheral blood lymphocytes.

### External Database Links

#### **UniProt:**

P01732 Related reagents
P10966 Related reagents

#### **Entrez Gene:**

925 CD8A Related reagents926 CD8B Related reagents

#### **Synonyms**

CD8B1, MAL

### **Fusion Partners**

Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63.653 myeloma cell line.

### **Specificity**

**Mouse anti Human CD8 antibody, clone LT8** recognizes the human CD8 cell surface glycoprotein expressed by a subset of peripheral blood T cells which express cytotoxic/suppressor activity. It is also expressed weakly on NK cells.

The CD8 antigen is a co-receptor for MHC Class I in conjunction with the T cell receptor, and is important in the selection process of CD8+ MHC Class I restricted T cells.

#### Flow Cytometry

Use 5ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

### References

- 1. Zarkesh-Esfahani, H. *et al.* (2001) High-dose leptin activates human leukocytes via receptor expression on monocytes. J Immunol. 167 (8): 4593-9.
- 2. Manninen, A. & Saksela, K. (2002) HIV-1 Nef interacts with inositol trisphosphate receptor to activate calcium signaling in T cells. <u>J Exp Med. 195 (8): 1023-32.</u>
- 3. Parnes, J.R. (1989) Molecular biology and function of CD4 and CD8. <u>Adv Immunol. 44:</u> <u>265-311.</u>
- 4. Kap, Y.S. *et al.* (2009) A monoclonal antibody selection for immunohistochemical examination of lymphoid tissues from non-human primates. <u>J Histochem Cytochem. 57:</u> 1159-67.
- 5. Hovden, A.O. *et al.* (2011) Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. <u>BMC Immunol. 12: 2.</u>
- 6. Nelson, M. *et al.* (2010) Characterization of lethal inhalational infection with *Francisella tularensis* in the common marmoset (*Callithrix jacchus*). <u>J Med Microbiol</u>. 59: 1107-13.
- 7. Gibbings, D.J. *et al.* (2007) CD8 alpha is expressed by human monocytes and enhances Fc gamma R-dependent responses. <u>BMC Immunol</u>. 8: 12.
- 8. Junker, A. et al. (2007) Multiple sclerosis: T-cell receptor expression in distinct brain

regions. Brain. 130: 2789-99.

- 9. Held, K. *et al.* (2011) Expression of herpes simplex virus 1-encoded microRNAs in human trigeminal ganglia and their relation to local T-cell infiltrates. <u>J Virol. 85 (19):</u> 9680-5.
- 10. Hood SP *et al.* (2014) Changes in immune cell populations in the periphery and liver of GBV-B-infected and convalescent tamarins (*Saguinus labiatus*). <u>Virus Res. 179:</u> 93-101.
- 11. Nelson, M. & Loveday, M. (2014) Exploring the innate immunological response of an alternative nonhuman primate model of infectious disease; the common marmoset. <u>J. Immunol Res. 2014</u>: 913632.
- 12. Manivannan, K. *et al.* (2016) CADM1/TSLC1 Identifies HTLV-1-Infected Cells and Determines Their Susceptibility to CTL-Mediated Lysis. <u>PLoS Pathog. 12 (4): e1005560.</u>
- 13. Gross, C.C. *et al.* (2016) Impaired NK-mediated regulation of T-cell activity in multiple sclerosis is reconstituted by IL-2 receptor modulation. <u>Proc Natl Acad Sci U S A. 113 (21):</u> E2973-82.
- 14. Dunham, J. *et al.* (2016) Blockade of CD127 Exerts a Dichotomous Clinical Effect in Marmoset Experimental Autoimmune Encephalomyelitis. <u>J Neuroimmune Pharmacol. 11</u> (1): 73-83.
- 15. Bughani, U. *et al.* (2017) T cell activation and differentiation is modulated by a CD6 domain 1 antibody Itolizumab. <u>PLoS One. 12 (7): e0180088.</u>
- 16. Philippens, I.H. *et al.* (2017) Acceleration of Amyloidosis by Inflammation in the Amyloid-Beta Marmoset Monkey Model of Alzheimer's Disease. <u>J Alzheimers Dis. 55 (1):</u> 101-113.

| Storage                          | Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.   |        |
|----------------------------------|--|--------|
| Guarantee                        | 12 months from date of despatch  |        |
| Acknowledgements                 | This product is covered by U.S. Patent No. 10,150,841 and related U.S. and f counterparts  | oreign |
| Health And Safety<br>Information | Material Safety Datasheet documentation #20471 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1226SBUV795">https://www.bio-rad-antibodies.com/SDS/MCA1226SBUV795</a> 20471 |        |
| Regulatory                       | For research purposes only   |        |

### Related Products

# **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody sales us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody sales de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

# Printed on 16 Oct 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint