Datasheet: MCA1226SBV515 BATCH NUMBER 100007653

Description:	MOUSE ANTI HUMAN CD8:StarBright Violet 515
Specificity:	CD8
Format:	StarBright Violet 515
Product Type:	Monoclonal Antibody
Clone:	LT8
lsotype:	lgG1
Quantity:	100 TESTS/0.5ml

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further					
	information. For gener	•				
	rad-antibodies.com/protocols.					
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry	-			Neat	
	Where this product has not been tested for use in a particular technique this does not					
	necessarily exclude its a guide only. It is reco system using appropri	mmended that	the use	er titrates the product f	g dilutions are given as or use in their own	
Target Species	Human					
Species Cross Reactivity	Reacts with: Marmoset, Chimpanzee, Cynomolgus monkey, Red-bellied Tamarin N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.					
Product Form	Purified IgG conjugated to StarBright Violet 515 - liquid					
Max Ex/Em	Fluorophore	Excitation Ma	x (nm)	Emission Max (nm)		
	StarBright Violet 515	402	()	516		
Preparation	Purified IgG prepared	by ion exchan	ge chro	matography from ascit	tes	
Buffer Solution	Phosphate buffered sa	aline				

Preservative Stabilisers	0.09% sodium azide (NaN ₃) 1% bovine serum albumin 0.1% Pluronic F68 0.1% PEG 3350 0.05% Tween 20
Immunogen	Normal human peripheral blood lymphocytes.
External Database Links	UniProt: <u>P01732</u> <u>Related reagents</u> <u>P10966</u> <u>Related reagents</u> Entrez Gene:
	925 CD8A <u>Related reagents</u> 926 CD8B <u>Related reagents</u>
Superume	
Synonyms	CD8B1, MAL
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63.653 myeloma cell line.
Specificity	 Mouse anti Human CD8 antibody, clone LT8 recognizes the human CD8 cell surface glycoprotein expressed by a subset of peripheral blood T cells which express cytotoxic/suppressor activity. It is also expressed weakly on NK cells. The CD8 antigen is a co-receptor for MHC Class I in conjunction with the T cell receptor, and is important in the selection process of CD8+ MHC Class I restricted T cells.
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 ⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
References	 Zarkesh-Esfahani, H. <i>et al.</i> (2001) High-dose leptin activates human leukocytes via receptor expression on monocytes. J Immunol. 167 (8): 4593-9. Manninen, A. & Saksela, K. (2002) HIV-1 Nef interacts with inositol trisphosphate receptor to activate calcium signaling in T cells. J Exp Med. 195 (8): 1023-32. Parnes, J.R. (1989) Molecular biology and function of CD4 and CD8. Adv Immunol. 44: 265-311. Kap, Y.S. <i>et al.</i> (2009) A monoclonal antibody selection for immunohistochemical examination of lymphoid tissues from non-human primates. J Histochem Cytochem. 57: 1159-67. Hovden, A.O. <i>et al.</i> (2011) Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. BMC Immunol. 12: 2. Nelson, M. <i>et al.</i> (2010) Characterization of lethal inhalational infection with <i>Francisella tularensis</i> in the common marmoset (<i>Callithrix jacchus</i>). J Med Microbiol. 59: 1107-13. Gibbings, D.J. <i>et al.</i> (2007) CD8 alpha is expressed by human monocytes and enhances Fc gamma R-dependent responses. BMC Immunol. 8: 12. Junker, A. <i>et al.</i> (2007) Multiple sclerosis: T-cell receptor expression in distinct brain

	 regions. <u>Brain. 130: 2789-99.</u> 9. Held, K. <i>et al.</i> (2011) Expression of herpes simplex virus 1-encoded microRN human trigeminal ganglia and their relation to local T-cell infiltrates. <u>J Virol. 85 (9680-5.</u> 10. Hood SP <i>et al.</i> (2014) Changes in immune cell populations in the periphery of GBV-B-infected and convalescent tamarins (<i>Saguinus labiatus</i>). <u>Virus Res. 1</u>: 93-101. 11. Nelson, M. & Loveday, M. (2014) Exploring the innate immunological respor alternative nonhuman primate model of infectious disease; the common marmoral <u>Immunol Res. 2014: 913632.</u> 12. Manivannan, K. <i>et al.</i> (2016) CADM1/TSLC1 Identifies HTLV-1-Infected Cell Determines Their Susceptibility to CTL-Mediated Lysis. <u>PLoS Pathog. 12 (4): e1</u> 13. Gross, C.C. <i>et al.</i> (2016) Impaired NK-mediated regulation of T-cell activity is sclerosis is reconstituted by IL-2 receptor modulation. <u>Proc Natl Acad Sci U S A E2973-82.</u> 14. Dunham, J. <i>et al.</i> (2016) Blockade of CD127 Exerts a Dichotomous Clinical Marmoset Experimental Autoimmune Encephalomyelitis. <u>J Neuroimmune Pharr</u> (1): 73-83. 15. Bughani, U. <i>et al.</i> (2017) T cell activation and differentiation is modulated by domain 1 antibody Itolizumab. <u>PLoS One. 12 (7): e0180088.</u> 16. Philippens, I.H. <i>et al.</i> (2017) Acceleration of Amyloidosis by Inflammation in Amyloid-Beta Marmoset Monkey Model of Alzheimer's Disease. <u>J Alzheimers D 101-113.</u> 	$\frac{19):}{2}$ and liver $\frac{79:}{79:}$ anse of an set. <u>J</u> and
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.	
Guarantee	12 months from date of despatch	
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and for counterparts	eign
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA1226SBV515 20471	
Regulatory	For research purposes only	

Related Products

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide
America	Fax: +1 919 878 3751	
	Email: antibody_sales_us@bio-rad.c	com

Tel: +44 (0)1865 852 700 **Europe** Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint