

Datasheet: MCA1227F

| | |
|----------------------|-----------------------------|
| Description: | MOUSE ANTI HUMAN CD42a:FITC |
| Specificity: | CD42a |
| Other names: | GPIX |
| Format: | FITC |
| Product Type: | Monoclonal Antibody |
| Clone: | GRP-P |
| Isotype: | IgG1 |
| Quantity: | 0.1 mg |

Product Details

RRID AB_321698

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ■ | | | Neat |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Human

Species Cross Reactivity Reacts with: Mink, Dog
N.B. Antibody reactivity and working conditions may vary between species.

Product Form Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid.

| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) |
|-----------|-------------|---------------------|-------------------|
| | FITC | 490 | 525 |

Preparation Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative Stabilisers 0.09% Sodium Azide
1% Bovine Serum Albumin

Approx. Protein Concentrations IgG concentration 0.1 mg/ml

Immunogen Human red blood cells and platelets.

| | |
|--------------------------------|---|
| External Database Links | UniProt: P14770 Related reagents Entrez Gene: 2815 GP9 Related reagents |
| Fusion Partners | Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line. |
| Specificity | <p>Mouse anti Human CD42a antibody, clone GRP-P recognizes the platelet GPIX glycoprotein, a 23kDa surface marker expressed by platelets and megakaryocytes. Platelet GPIX is also known as CD42a.</p> <p>The CD42 complex is the major platelet receptor for von Willebrand factor.</p> |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood |
| References | <ol style="list-style-type: none"> 1. Sarma, J. <i>et al.</i> (2002) Increased platelet binding to circulating monocytes in acute coronary syndromes. Circulation. 105 (18): 2166-71. 2. Bournazos, S. <i>et al.</i> (2008) Monocyte functional responsiveness after PSGL-1-mediated platelet adhesion is dependent on platelet activation status. Arterioscler Thromb Vasc Biol. 28: 1491-8. 3. Aasted, B. <i>et al.</i> (2007) Reactivity of monoclonal antibodies to human CD antigens with cells from mink. Vet Immunol Immunopathol. 119: 27-37. 4. Perdomo, J. <i>et al.</i> (2011) Quinine-induced thrombocytopenia: drug-dependent GPIb/IX antibodies inhibit megakaryocyte and proplatelet production in vitro. Blood. 117: 5975-86. 5. Harding, S.A. <i>et al.</i> (2004) Increased CD40 ligand and platelet-monocyte aggregates in patients with type 1 diabetes mellitus. Atherosclerosis. 176: 321-5. 6. Harding, S.A. <i>et al.</i> (2007) Clopidogrel reduces platelet-leucocyte aggregation, monocyte activation and RANTES secretion in type 2 diabetes mellitus. Heart. 92: 1335-7. 7. Eisbacher, M. <i>et al.</i> (2001) Inducible expression of the megakaryocyte-specific gene glycoprotein IX is mediated through an Ets binding site and involves upstream activation of extracellular signal-regulated kinase. Cell Growth Differ. 12: 435-45. 8. Vettore, S. <i>et al.</i> (2008) Novel point mutation in a leucine-rich repeat of the GPIbalpha chain of the platelet von Willebrand factor receptor, GPIb/IX/V, resulting in an inherited dominant form of Bernard-Soulier syndrome affecting two unrelated families: the N41H variant. Haematologica. 93: 1743-7. 9. Fox, S.C. <i>et al.</i> (2004) Quantitation of platelet aggregation and microaggregate formation in whole blood by flow cytometry. Platelets. 15: 85-93. 10. Din, J.N. <i>et al.</i> (2010) Effect of moderate walnut consumption on lipid profile, arterial stiffness and platelet activation in humans. Eur J Clin Nutr. 2011 Feb;65(2):234-9. 11. Din, J.N. <i>et al.</i> (2008) Dietary intervention with oil rich fish reduces platelet-monocyte aggregation in man. Atherosclerosis. 197: 290-6. 12. Schuberth, H.J. <i>et al.</i> (2007) Reactivity of cross-reacting monoclonal antibodies with canine leukocytes, platelets and erythrocytes. Vet Immunol Immunopathol. 119: 47-55. |
| Storage | <p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.</p> |

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

| | |
|------------------|---------------------------------|
| Guarantee | 18 months from date of despatch |
|------------------|---------------------------------|

| | |
|--------------------------------------|--|
| Health And Safety Information | Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf |
|--------------------------------------|--|

| | |
|-------------------|----------------------------|
| Regulatory | For research purposes only |
|-------------------|----------------------------|

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M344828:190118'

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)