

## Datasheet: MCA1227FT

<b>Description:</b>	MOUSE ANTI HUMAN CD42a:FITC
<b>Specificity:</b>	CD42a
<b>Other names:</b>	GPIX
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	GRP-P
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	25 µg

## Product Details

**RRID** AB\_2263686

**Applications** This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**Target Species** Human

**Species Cross Reactivity** Reacts with: Mink, Dog  
**N.B.** Antibody reactivity and working conditions may vary between species.

**Product Form** Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid.

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

**Preparation** Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

**Buffer Solution** Phosphate buffered saline

**Preservative Stabilisers** 0.09% Sodium Azide  
1% Bovine Serum Albumin

**Approx. Protein Concentrations** IgG concentration 0.1 mg/ml

**Immunogen** Human red blood cells and platelets.

<b>External Database Links</b>	<b>UniProt:</b> <a href="#">P14770</a> <a href="#">Related reagents</a>  <b>Entrez Gene:</b> <a href="#">2815</a> GP9 <a href="#">Related reagents</a>
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Human CD42a antibody, clone GRP-P</b> recognizes the platelet GPIX glycoprotein, a 23kDa surface marker expressed by platelets and megakaryocytes. Platelet GPIX is also known as CD42a.</p> <p>The CD42 complex is the major platelet receptor for von Willebrand factor.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells or 100ul whole blood
<b>References</b>	<ol style="list-style-type: none"> <li>Sarma, J. <i>et al.</i> (2002) Increased platelet binding to circulating monocytes in acute coronary syndromes. <a href="#">Circulation. 105 (18): 2166-71.</a></li> <li>Bournazos, S. <i>et al.</i> (2008) Monocyte functional responsiveness after PSGL-1-mediated platelet adhesion is dependent on platelet activation status. <a href="#">Arterioscler Thromb Vasc Biol. 28: 1491-8.</a></li> <li>Aasted, B. <i>et al.</i> (2007) Reactivity of monoclonal antibodies to human CD antigens with cells from mink. <a href="#">Vet Immunol Immunopathol. 119: 27-37.</a></li> <li>Perdomo, J. <i>et al.</i> (2011) Quinine-induced thrombocytopenia: drug-dependent GPIb/IX antibodies inhibit megakaryocyte and proplatelet production in vitro. <a href="#">Blood. 117: 5975-86.</a></li> <li>Harding, S.A. <i>et al.</i> (2004) Increased CD40 ligand and platelet-monocyte aggregates in patients with type 1 diabetes mellitus. <a href="#">Atherosclerosis. 176: 321-5.</a></li> <li>Harding, S.A. <i>et al.</i> (2007) Clopidogrel reduces platelet-leucocyte aggregation, monocyte activation and RANTES secretion in type 2 diabetes mellitus. <a href="#">Heart. 92: 1335-7.</a></li> <li>Eisbacher, M. <i>et al.</i> (2001) Inducible expression of the megakaryocyte-specific gene glycoprotein IX is mediated through an Ets binding site and involves upstream activation of extracellular signal-regulated kinase. <a href="#">Cell Growth Differ. 12: 435-45.</a></li> <li>Vettore, S. <i>et al.</i> (2008) Novel point mutation in a leucine-rich repeat of the GPIIb/IIIa chain of the platelet von Willebrand factor receptor, GPIIb/IIIa/V, resulting in an inherited dominant form of Bernard-Soulier syndrome affecting two unrelated families: the N41H variant. <a href="#">Haematologica. 93: 1743-7.</a></li> <li>Fox, S.C. <i>et al.</i> (2004) Quantitation of platelet aggregation and microaggregate formation in whole blood by flow cytometry. <a href="#">Platelets. 15: 85-93.</a></li> <li>Din, J.N. <i>et al.</i> (2010) Effect of moderate walnut consumption on lipid profile, arterial stiffness and platelet activation in humans. <a href="#">Eur J Clin Nutr. 2011 Feb;65(2):234-9.</a></li> <li>Din, J.N. <i>et al.</i> (2008) Dietary intervention with oil rich fish reduces platelet-monocyte aggregation in man. <a href="#">Atherosclerosis. 197: 290-6.</a></li> <li>Schuberth, H.J. <i>et al.</i> (2007) Reactivity of cross-reacting monoclonal antibodies with canine leukocytes, platelets and erythrocytes. <a href="#">Vet Immunol Immunopathol. 119: 47-55.</a></li> </ol>
<b>Storage</b>	<p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.</p>

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

<b>Guarantee</b>	18 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10041 available at: 10041: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf</a>
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M344841:190118'

**Printed on 20 May 2019**

---

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)