

Datasheet: MCA1227PE

Description:	MOUSE ANTI HUMAN CD42a:RPE
Specificity:	CD42a
Other names:	GPIX
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	GRP-P
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Mink, Dog

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized

Reconstitution

Reconstitute with 1 ml distilled water

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
RPE 488nm laser	496	578

Preparation

Purified IgG prepared by affinity chromatography on Protein A

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin 5% Sucrose
Immunogen	Human red blood cells and platelets.
External Database Links	<p>UniProt: P14770 Related reagents</p> <p>Entrez Gene: 2815 GP9 Related reagents</p>
RRID	AB_321699
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
Specificity	<p>Mouse anti Human CD42a antibody, clone GRP-P recognizes the platelet GPIX glycoprotein, a 23kDa surface marker expressed by platelets and megakaryocytes. Platelet GPIX is also known as CD42a.</p> <p>The CD42 complex is the major platelet receptor for von Willebrand factor.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood
References	<ol style="list-style-type: none"> Sarma, J. <i>et al.</i> (2002) Increased platelet binding to circulating monocytes in acute coronary syndromes. Circulation. 105 (18): 2166-71. Bournazos, S. <i>et al.</i> (2008) Monocyte functional responsiveness after PSGL-1-mediated platelet adhesion is dependent on platelet activation status. Arterioscler Thromb Vasc Biol. 28: 1491-8. Aasted, B. <i>et al.</i> (2007) Reactivity of monoclonal antibodies to human CD antigens with cells from mink. Vet Immunol Immunopathol. 119: 27-37. Perdomo, J. <i>et al.</i> (2011) Quinine-induced thrombocytopenia: drug-dependent GPIb/IX antibodies inhibit megakaryocyte and proplatelet production in vitro. Blood. 117: 5975-86. Harding, S.A. <i>et al.</i> (2004) Increased CD40 ligand and platelet-monocyte aggregates in patients with type 1 diabetes mellitus. Atherosclerosis. 176: 321-5. Harding, S.A. <i>et al.</i> (2007) Clopidogrel reduces platelet-leucocyte aggregation, monocyte activation and RANTES secretion in type 2 diabetes mellitus. Heart. 92: 1335-7. Eisbacher, M. <i>et al.</i> (2001) Inducible expression of the megakaryocyte-specific gene glycoprotein IX is mediated through an Ets binding site and involves upstream activation of extracellular signal-regulated kinase. Cell Growth Differ. 12: 435-45. Vettore, S. <i>et al.</i> (2008) Novel point mutation in a leucine-rich repeat of the GPIbalpha chain of the platelet von Willebrand factor receptor, GPIb/IX/V, resulting in an inherited dominant form of Bernard-Soulier syndrome affecting two unrelated families: the N41H variant. Haematologica. 93: 1743-7.

9. Fox, S.C. *et al.* (2004) Quantitation of platelet aggregation and microaggregate formation in whole blood by flow cytometry. [Platelets. 15: 85-93.](#)
10. Din, J.N. *et al.* (2010) Effect of moderate walnut consumption on lipid profile, arterial stiffness and platelet activation in humans. [Eur J Clin Nutr. 2011 Feb;65\(2\):234-9.](#)
11. Din, J.N. *et al.* (2008) Dietary intervention with oil rich fish reduces platelet-monocyte aggregation in man. [Atherosclerosis. 197: 290-6.](#)
12. Schuberth, H.J. *et al.* (2007) Reactivity of cross-reacting monoclonal antibodies with canine leukocytes, platelets and erythrocytes. [Vet Immunol Immunopathol. 119: 47-55.](#)
13. Tunströmer, K. *et al.* (2018) Quantification of Platelet Contractile Movements during Thrombus Formation. [Thromb Haemost. 118 \(9\): 1600-11.](#)

Storage Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #20487 available at: 20487: <https://www.bio-rad-antibodies.com/uploads/MSDS/20487.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M375293:210104'

Printed on 22 Mar 2021