

Datasheet: MCA1258PECY7

Description:	RAT ANTI MOUSE CD45R:RPE-Cy7
Specificity:	CD45R
Other names:	B220, LY-5
Format:	RPE-CY7
Product Type:	Monoclonal Antibody
Clone:	RA3-6B2
Isotype:	IgG2a
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			neat - 1/5

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Mouse

Species Cross Reactivity

Reacts with: Human, Cat

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to R. Phycoerythrin -Cy7 (RPE-Cy7) - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
RPE-Cy7 488nm laser	496	695

Preparation

Purified IgG prepared by affinity chromatography on Protein A

Buffer Solution

Phosphate buffered saline

Preservative

0.1% Sodium Azide (NaN₃)

Stabilisers	Stabilizing agent
Immunogen	Murine leukemia-induced pre-B tumor cells (RAW112)
External Database Links	<p>UniProt: P06800 Related reagents</p> <p>Entrez Gene: 19264 Ptprc Related reagents</p>
Synonyms	Ly-5
RRID	AB_905963
Fusion Partners	Spleen cells from immunized Lewis rats were fused with cells of the rat S194/5 XX0.BU-1 myeloma cell line
Specificity	<p>Rat anti Mouse CD45R antibody, clone RA3-6B2 recognizes murine CD45R, a form of the CD45 antigen expressed by B cells and lytically active subsets of NK cells and non-MHC restricted CTL's. Rat anti Mouse CD45R antibody, clone RA3-6B2 immunoprecipitates the high molecular weight form of CD45 (220 kDa).</p> <p>Rat anti Mouse CD45R antibody, clone RA3-6B2 is suitable for plp fixed paraffin embedded tissues (Whiteland et al.1995).</p>
Flow Cytometry	<p>Use 10 ul of the suggested working dilution to label 10^6 cells in 100ul.</p> <p>The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B).</p>
References	<ol style="list-style-type: none"> Coffman, R.L. (1982) Surface antigen expression and immunoglobulin gene rearrangement during mouse pre-B cell development. Immunol Rev. 69: 5-23. Rosmalen, J.G. et al. (2000) Subsets of macrophages and dendritic cells in nonobese diabetic mouse pancreatic inflammatory infiltrates: correlation with the development of diabetes. Lab Invest. 80 (1): 23-30. Whiteland, J.L. et al. (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. J Histochem Cytochem. 43 (3): 313-20. Spangrude, G.J. et al. (1988) Purification and characterization of mouse hematopoietic stem cells. Science. 241: 58-62. Spangrude, G.J. et al. (1988) Two rare populations of mouse Thy-1lo bone marrow cells repopulate the thymus. J Exp Med. 167 (5): 1671-83. Holmes, K.L. et al. (1986) Analysis of neoplasms induced by Cas-Br-M MuLV tumor extracts. J Immunol. 137 (2): 679-88. Ankeny, D.P. et al. (2009) B cells produce pathogenic antibodies and impair recovery after spinal cord injury in mice. J Clin Invest. 119: 2990-9. Lundqvist, J. et al. (2010) Concomitant infection decreases the malaria burden but

- escalates relapsing fever borreliosis. [Infect Immun. 78 \(5\): 1924-30.](#)
9. Herrmann, I. *et al.* (2006) Streptococcus pneumoniae Infection aggravates experimental autoimmune encephalomyelitis via Toll-like receptor 2. [Infect Immun. 74: 4841-8.](#)
10. Kleiter, I. *et al.* (2010) Smad7 in T cells drives T helper 1 responses in multiple sclerosis and experimental autoimmune encephalomyelitis. [Brain. 133: 1067-81.](#)
11. Lacroix-Lamande, S. *et al.* (2009) Neonate intestinal immune response to CpG oligodeoxynucleotide stimulation. [PLoS One. 4: e8291.](#)
12. Bertilaccio, M.T. *et al.* (2011) Lack of TIR8/SIGIRR triggers progression of chronic lymphocytic leukemia in mouse models. [Blood. 118: 660-9.](#)
13. Gengozian, N. *et al.* (2005) Characterization of a monoclonal antibody identifying a CD45RA antigen on feline leukocytes. [Vet Immunol Immunopathol. 108: 253-64.](#)
14. Giuriato, S. *et al.* (2010) Conditional TPM3-ALK and NPM-ALK transgenic mice develop reversible ALK-positive early B-cell lymphoma/leukemia. [Blood. 115: 4061-70.](#)
15. Hawke, S. *et al.* (1998) Long-term persistence of activated cytotoxic T lymphocytes after viral infection of the central nervous system. [J Exp Med. 187: 1575-82.](#)
16. Nakaya, T. *et al.* (2010) Critical role of Pcid2 in B cell survival through the regulation of MAD2 expression. [J Immunol. 185: 5180-7.](#)
17. Perry, M.J. *et al.* (2000) Effects of high-dose estrogen on murine hematopoietic bone marrow precede those on osteogenesis. [Am J Physiol Endocrinol Metab. 279: E1159-65.](#)
18. Gibson-Corley, K.N. *et al.* (2016) A method for histopathological study of the multifocal nature of spinal cord lesions in murine experimental autoimmune encephalomyelitis. [PeerJ. 4: e1600.](#)
19. Soejima, M. *et al.* (2011) Role of innate immunity in a murine model of histidyl-transfer RNA synthetase (Jo-1)-mediated myositis. [Arthritis Rheum. 63: 479-87.](#)
20. Stevenson, P.G. *et al.* (2002) Uncoupling of virus-induced inflammation and anti-viral immunity in the brain parenchyma. [J Gen Virol. 83: 1735-43.](#)
21. Fanning, S. *et al.* (2012) Bifidobacterial surface-exopolysaccharide facilitates commensal-host interaction through immune modulation and pathogen protection. [Proc Natl Acad Sci U S A. 109 \(6\): 2108-13.](#)
22. Ruf, M.T. *et al.* (2012) Chemotherapy-Associated Changes of Histopathological Features of Mycobacterium ulcerans Lesions in a Buruli Ulcer Mouse Model. [Antimicrob Agents Chemother. 56: 687-96.](#)
23. Carpenter, R.S. *et al.* (2015) Traumatic spinal cord injury in mice with human immune systems. [Exp Neurol. 271: 432-44.](#)
24. Lastrucci, C. *et al.* (2015) Molecular and cellular profiles of the resolution phase in a damage-associated molecular pattern (DAMP)-mediated peritonitis model and revelation of leukocyte persistence in peritoneal tissues. [FASEB J. 29 \(5\): 1914-29.](#)

Storage

Store at +4°C.

DO NOT FREEZE

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

Guaranteed until date of expiry. Please see product label.

Acknowledgements Cy® and CyDye® are registered trademarks of GE Healthcare

Health And Safety Information Material Safety Datasheet documentation #10331 available at:
10331: <https://www.bio-rad-antibodies.com/uploads/MSDS/10331.pdf>

Regulatory For research purposes only

Related Products

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M350378:190307'

Printed on 16 Oct 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)