

# Datasheet: MCA1260G BATCH NUMBER 150150

Description:	RAT ANTI MOUSE CD25
Specificity:	CD25
Other names:	IL-2R ALPHA CHAIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	PC61.5.3
Isotype:	lgG1

0.25 mg

## **Product Details**

#### **Applications**

Quantity:

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			
Immunohistology - Frozen	•			
Immunohistology - Paraffin			•	
ELISA			•	
Immunoprecipitation	•			
Western Blotting			•	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein C supernatant	G from tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> )	

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	B6.1 CTL cell line.
External Database Links	UniProt: P01590 Related reagents  Entrez Gene: 16184 Il2ra Related reagents
Synonyms	ll2r
RRID	AB_2125347
Fusion Partners	Spleen cells from immunized OFA rats were fuzed with cells of the P3X63Ag8.653 mouse myeloma cell line.
Specificity	Rat anti Mouse CD25 antibody, clone PC61.5.3 reacts with the low affinity alpha chain of the interleukin-2 receptor present on activated T and B cells in mice. Rat anti Mouse CD25 antibody, clone PC61.5.3 is reported to inhibit IL-2 binding and IL-2 dependent proliferation.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1 x $10^6$ cells in 100ul.
References	<ol> <li>Moreau, J.L. <i>et al.</i> (1987) Monoclonal antibodies identify three epitope clusters on the mouse p55 subunit of the interleukin 2 receptor: relationship to the interleukin 2-binding site. Eur J Immunol. 17 (7): 929-35.</li> <li>Hashimoto, N. <i>et al.</i> (1986) Dissociation of interleukin 2-dependent and -independent B cell proliferation with monoclonal anti-interleukin 2 receptor antibody. Eur J Immunol. 16 (3): 317-20.</li> <li>Lowenthal, J.W. <i>et al.</i> (1985) High and low affinity IL 2 receptors: analysis by IL 2 dissociation rate and reactivity with monoclonal anti-receptor antibody PC61. J Immunol. 135 (6): 3988-94.</li> <li>Ceredig, R. <i>et al.</i> (1985) Expression of interleukin-2 receptor as a differentiation marker on intrathymic stem cells. Nature 314: 98-100.</li> <li>Yaqoob, P. &amp; Calder, P.C. (1997) Glutamine requirement of proliferating T lymphocytes. Nutrition. 13 (7-8): 646-51.</li> <li>Scotland, R.S. <i>et al.</i> (2011) Sex differences in resident immune cell phenotype underlie more efficient acute inflammatory responses in female mice. Blood. 118 (22): 5918-27.</li> <li>Karali, D. <i>et al.</i> (2016) T cell regulation by <i>Phlomis lanata</i> protein extracts in mice. Pharm Biol. 54 (2): 207-14.</li> <li>Szuster-Ciesielska, A. <i>et al.</i> (2019) Immunogenic Evaluation of Ribosomal P-Protein Antigen P0, P1, and P2 and Pentameric Protein Complex P0-(P1-P2)<sub>2</sub> of <i>Plasmodium falciparum</i> in a Mouse Model. J Immunol Res. 2019: 9264217.</li> </ol>

Storage Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may

denature the antibody. Should this product contain a precipitate we recommend

microcentrifugation before use.

**Guarantee** 12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at:

https://www.bio-rad-antibodies.com/SDS/MCA1260G

10040

**Regulatory** For research purposes only

#### Related Products

### **Recommended Secondary Antibodies**

Goat Anti Rat IgG (STAR73...) RPE

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) DyLight®550, DyLight®650, DyLight®800

Rabbit Anti Rat IgG (STAR21...) HRP

Rabbit Anti Rat IgG (STAR16...) <u>DyLight®800</u>
Goat Anti Rat IgG (STAR131...) <u>Alk. Phos., Biotin</u>

Rabbit Anti Rat IgG (STAR17...) FITC

Goat Anti Rat IgG (STAR72...) HRP

Goat Anti Rat IgG (STAR69...) FITC

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Fax: +44 (0)1865 852 739
Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M365018:200529'

#### Printed on 07 Sep 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint