

Datasheet: MCA1260SBUV605

Description:	RAT ANTI MOUSE CD25:StarBright UltraViolet 605
Specificity:	CD25
Other names:	IL-2R ALPHA CHAIN
Format:	StarBright UltraViolet 605
Product Type:	Monoclonal Antibody
Clone:	PC61.5.3
Isotype:	lgG1
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse				
Product Form	Purified IgG conjugated to StarBright UltraViolet 605 - liquid				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm		
	StarBright UltraViolet 605	340	609		
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue consupernatant				
Buffer Solution	Phosphate buffered saline				
Preservative Stabilisers	0.09% sodium azide (NaN ₃) 1% bovine serum albumin 0.1% Pluronic F68 0.1% PEG 3350				

B6.1 CTL cell line.

External Database Links

UniProt:

P01590 Related reagents

Entrez Gene:

16184 Il2ra Related reagents

Synonyms

II2r

Fusion Partners

Spleen cells from immunized OFA rats were fused with cells of the P3X63Ag8.653 mouse myeloma cell line.

Specificity

Rat anti Mouse CD25 antibody, clone PC61.5.3 reacts with the low affinity alpha chain of the interleukin-2 receptor present on activated T and B cells in mice. Rat anti Mouse CD25 antibody, clone PC61.5.3 is reported to inhibit IL-2 binding and IL-2 dependent proliferation.

Flow Cytometry

Use 5μ I of the suggested working dilution to label 10^6 cells in 100μ I. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

References

- 1. Ceredig, R. *et al.* (1985) Expression of interleukin-2 receptor as a differentiation marker on intrathymic stem cells. Nature 314: 98-100.
- 2. Lowenthal, J.W. *et al.* (1985) High and low affinity IL 2 receptors: analysis by IL 2 dissociation rate and reactivity with monoclonal anti-receptor antibody PC61. <u>J Immunol.</u> 135 (6): 3988-94.
- 3. Hashimoto, N. *et al.* (1986) Dissociation of interleukin 2-dependent and -independent B cell proliferation with monoclonal anti-interleukin 2 receptor antibody. <u>Eur J Immunol. 16</u> (3): 317-20.
- 4. Moreau, J.L. *et al.* (1987) Monoclonal antibodies identify three epitope clusters on the mouse p55 subunit of the interleukin 2 receptor: relationship to the interleukin 2-binding site. Eur J Immunol. 17 (7): 929-35.
- 5. Yaqoob, P. & Calder, P.C. (1997) Glutamine requirement of proliferating T lymphocytes. Nutrition. 13 (7-8): 646-51.
- 6. Scotland, R.S. *et al.* (2011) Sex differences in resident immune cell phenotype underlie more efficient acute inflammatory responses in female mice. <u>Blood. 118 (22): 5918-27.</u>
- 7. Karali, D. *et al.* (2016) T cell regulation by *Phlomis lanata* protein extracts in mice. Pharm Biol. 54 (2): 207-14.
- 8. Szuster-Ciesielska, A. *et al.* (2019) Immunogenic Evaluation of Ribosomal P-Protein Antigen P0, P1, and P2 and Pentameric Protein Complex P0-(P1-P2)₂ of *Plasmodium falciparum* in a Mouse Model. J Immunol Res. 2019: 9264217.
- 9. Curina, G. *et al.* (2018) Evaluation of immune responses in mice and sheep inoculated with a live attenuated *Brucella melitensis*. REV1 vaccine produced in bioreactor. <u>Vet Immunol Immunopathol</u>. 198: 44-53.
- 10. Arad, T. et al. (2021) CD200 -dependent and -independent immune-modulatory

11. Roca, C.P. et al. (2023) A cross entropy test allows quantitative statistical comparison of t-SNE and UMAP representations Cell Reports Methods. 3 (1): 100390. Storage Store at +4°C. >DO NOT FREEZE. This product should be stored undiluted. Guarantee 12 months from date of despatch Acknowledgements This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts **Health And Safety** Material Safety Datasheet documentation #20471 available at: Information https://www.bio-rad-antibodies.com/SDS/MCA1260SBUV605 20471 Regulatory For research purposes only

functions of neural stem cells. Stem Cell Res. 56: 102559.

Related Products

Recommended Useful Reagents

MOUSE SEROBLOCK FCR (BUF041A)
MOUSE SEROBLOCK FCR (BUF041B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M408797:221014'

Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint