

Datasheet: MCA1260SBUV740

BATCH NUMBER 100007568

Description:	RAT ANTI MOUSE CD25:StarBright UltraViolet 740
Specificity:	CD25
Other names:	IL-2R ALPHA CHAIN
Format:	StarBright UltraViolet 740
Product Type:	Monoclonal Antibody
Clone:	PC61.5.3
Isotype:	IgG1
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to StarBright UltraViolet 740 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	StarBright UltraViolet 740	344	743
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin 0.1% Pluronic F68		

0.1% PEG 3350
0.05% Tween 20

Immunogen B6.1 CTL cell line.

External Database Links

UniProt:

[P01590](#) [Related reagents](#)

Entrez Gene:

[16184](#) Il2ra [Related reagents](#)

Synonyms Il2r

Fusion Partners Spleen cells from immunized OFA rats were fused with cells of the P3X63Ag8.653 mouse myeloma cell line.

Specificity **Rat anti Mouse CD25 antibody, clone PC61.5.3** reacts with the low affinity alpha chain of the interleukin-2 receptor present on activated T and B cells in mice. Rat anti Mouse CD25 antibody, clone PC61.5.3 is reported to inhibit IL-2 binding and IL-2 dependent proliferation.

Flow Cytometry Use 5µl of the suggested working dilution to label 10⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

References

1. Ceredig, R. *et al.* (1985) Expression of interleukin-2 receptor as a differentiation marker on intrathymic stem cells. [Nature 314: 98-100.](#)
2. Lowenthal, J.W. *et al.* (1985) High and low affinity IL 2 receptors: analysis by IL 2 dissociation rate and reactivity with monoclonal anti-receptor antibody PC61. [J Immunol. 135 \(6\): 3988-94.](#)
3. Hashimoto, N. *et al.* (1986) Dissociation of interleukin 2-dependent and -independent B cell proliferation with monoclonal anti-interleukin 2 receptor antibody. [Eur J Immunol. 16 \(3\): 317-20.](#)
4. Moreau, J.L. *et al.* (1987) Monoclonal antibodies identify three epitope clusters on the mouse p55 subunit of the interleukin 2 receptor: relationship to the interleukin 2-binding site. [Eur J Immunol. 17 \(7\): 929-35.](#)
5. Yaqoob, P. & Calder, P.C. (1997) Glutamine requirement of proliferating T lymphocytes. [Nutrition. 13 \(7-8\): 646-51.](#)
6. Scotland, R.S. *et al.* (2011) Sex differences in resident immune cell phenotype underlie more efficient acute inflammatory responses in female mice. [Blood. 118 \(22\): 5918-27.](#)
7. Karali, D. *et al.* (2016) T cell regulation by *Phlomis lanata* protein extracts in mice. [Pharm Biol. 54 \(2\): 207-14.](#)
8. Szuster-Ciesielska, A. *et al.* (2019) Immunogenic Evaluation of Ribosomal P-Protein Antigen P0, P1, and P2 and Pentameric Protein Complex P0-(P1-P2)₂ of *Plasmodium falciparum* in a Mouse Model. [J Immunol Res. 2019: 9264217.](#)
9. Curina, G. *et al.* (2018) Evaluation of immune responses in mice and sheep inoculated with a live attenuated *Brucella melitensis*. REV1 vaccine produced in bioreactor. [Vet Immunol Immunopathol. 198: 44-53.](#)

10. Arad, T. *et al.* (2021) CD200 -dependent and -independent immune-modulatory functions of neural stem cells. [Stem Cell Res. 56: 102559.](#)
11. Roca, C.P. *et al.* (2023) A cross entropy test allows quantitative statistical comparison of t-SNE and UMAP representations [Cell Reports Methods. 3 \(1\): 100390.](#)

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA1260SBUV740 20471
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M417322:230314'

Printed on 12 Dec 2024