

Datasheet: MCA1260SBV440 BATCH NUMBER 64601885

Description:	RAT ANTI MOUSE CD25:StarBright Violet 440			
Specificity:	CD25			
Other names:	IL-2R ALPHA CHAIN			
Format:	StarBright Violet 440			
Product Type:	Monoclonal Antibody			
Clone:	PC61.5.3			
Isotype:	lgG1			
Quantity:	100 TESTS/0.5ml			

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes No	Not Determined	Suggested Dilution		
	Flow Cytometry	•		Neat		
	Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.					
Target Species	Mouse					
Product Form	Purified IgG conjugated to StarBright Violet 440 - liquid					
Max Ex/Em	Fluorophore	Excitation Max (nm) Emission Max (nm)			
	StarBright Violet 440	383	436			
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant					
Buffer Solution	Phosphate buffered saline					
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin 0.1% Pluronic F68 0.1% PEG 3350					

Immunogen	B6.1 CTL cell line.
External Database Links	UniProt: <u>P01590</u> <u>Related reagents</u> Entrez Gene: <u>16184</u> II2ra <u>Related reagents</u>
Synonyms	ll2r
Fusion Partners	Spleen cells from immunized OFA rats were fused with cells of the P3X63Ag8.653 mouse myeloma cell line.
Specificity	Rat anti Mouse CD25 antibody, clone PC61.5.3 reacts with the low affinity alpha chain of the interleukin-2 receptor present on activated T and B cells in mice. Rat anti Mouse CD25 antibody, clone PC61.5.3 is reported to inhibit IL-2 binding and IL-2 dependent proliferation.
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 ⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
References	 Ceredig, R. <i>et al.</i> (1985) Expression of interleukin-2 receptor as a differentiation marker on intrathymic stem cells. <u>Nature 314: 98-100</u>. Lowenthal, J.W. <i>et al.</i> (1985) High and low affinity IL 2 receptors: analysis by IL 2 dissociation rate and reactivity with monoclonal anti-receptor antibody PC61. <u>J Immunol.</u> 135 (6): 3988-94. Hashimoto, N. <i>et al.</i> (1986) Dissociation of interleukin 2-dependent and -independent B cell proliferation with monoclonal anti-interleukin 2 receptor antibody. <u>Eur J Immunol. 16</u> (3): 317-20. Moreau, J.L. <i>et al.</i> (1987) Monoclonal antibodies identify three epitope clusters on the mouse p55 subunit of the interleukin 2 receptor: relationship to the interleukin 2-binding site. <u>Eur J Immunol. 17 (7): 929-35.</u> Yaqoob, P. & Calder, P.C. (1997) Glutamine requirement of proliferating T lymphocytes. <u>Nutrition. 13 (7-8): 646-51.</u> Scotland, R.S. <i>et al.</i> (2011) Sex differences in resident immune cell phenotype underlie more efficient acute inflammatory responses in female mice. <u>Blood. 118 (22): 5918-27.</u> 7. Karali, D. <i>et al.</i> (2016) T cell regulation by <i>Phlomis lanata</i> protein extracts in mice. <u>Pharm Biol. 54 (2): 207-14.</u> Szuster-Ciesielska, A. <i>et al.</i> (2019) Immunogenic Evaluation of Ribosomal P-Protein Antigen P0, P1, and P2 and Pentameric Protein Complex P0-(P1-P2)₂ of <i>Plasmodium falciparum</i> in a Mouse Model. J Immunol Res. 2019: 9264217. Curina, G. <i>et al.</i> (2018) Evaluation of immune responses in mice and sheep inoculated with a live attenuated <i>Brucella melitensis.</i> REV1 vaccine produced in bioreactor. <u>Vet Immunol Immunopathol. 198: 44-53.</u> Arad, T. <i>et al.</i> (2021) CD200 -dependent and -independent immune-modulatory functions of neural stem cells. <u>Stem Cell Res. 56: 102559.</u> Roca, C.P. <i>et al.</i> (2023) A cross entropy test allows quantitative statistical comparison

Storage	Store at +4°C. DO NOT FREEZE.
	This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety	Material Safety Datasheet documentation #20438 available at:
Information	https://www.bio-rad-antibodies.com/SDS/MCA1260SBV440
	20438
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

MOUSE SEROBLOCK FcR (BUF041A)
MOUSE SEROBLOCK FcR (BUF041B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio	-rad.com	Email: antibody_sales_uk@bio-	rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M417965:230420'

Printed on 15 May 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint