

Datasheet: MCA1260SBY575 BATCH NUMBER 100007269

Description:	RAT ANTI MOUSE CD25:StarBright Yellow 575
Specificity:	CD25
Other names:	IL-2R ALPHA CHAIN
Format:	StarBright Yellow 575
Product Type:	Monoclonal Antibody
Clone:	PC61.5.3
Isotype:	lgG1
Quantity:	100 TESTS/0.5ml

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .				
		Yes No	Not Determined	Suggested Dilution	
	Flow Cytometry	-		Neat	
	Where this product has necessarily exclude its a guide only. It is recor system using appropria	use in such procedu nmended that the use	res. Suggested working er titrates the product f	g dilutions are given as	
Target Species	Mouse				
Product Form	Purified IgG conjugated to StarBright Yellow 575 - liquid				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)		
	StarBright Yellow 575	548	579		
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant				
Buffer Solution	Phosphate buffered saline				
Preservative Stabilisers	0.09% Sodium Azide (1% Bovine Serum Albu 0.1% Pluronic F68 0.1% PEG 3350	0,			

	0.05% Tween 20
Immunogen	B6.1 CTL cell line.
External Database Links	UniProt: P01590 Related reagents Entrez Gene: <u>16184</u> II2ra <u>Related reagents</u>
Synonyms	ll2r
Fusion Partners	Spleen cells from immunized OFA rats were fused with cells of the P3X63Ag8.653 mouse myeloma cell line.
Specificity	Rat anti Mouse CD25 antibody, clone PC61.5.3 reacts with the low affinity alpha chain of the interleukin-2 receptor present on activated T and B cells in mice. Rat anti Mouse CD25 antibody, clone PC61.5.3 is reported to inhibit IL-2 binding and IL-2 dependent proliferation.
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 ⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
References	 Ceredig, R. <i>et al.</i> (1985) Expression of interleukin-2 receptor as a differentiation marker on intrathymic stem cells. <u>Nature 314: 98-100.</u> Lowenthal, J.W. <i>et al.</i> (1985) High and low affinity IL 2 receptors: analysis by IL 2 dissociation rate and reactivity with monoclonal anti-receptor antibody PC61. <u>J Immunol.</u> <u>135 (6): 3988-94.</u> Hashimoto, N. <i>et al.</i> (1986) Dissociation of interleukin 2-dependent and -independent B cell proliferation with monoclonal anti-interleukin 2 receptor antibody. <u>Eur J Immunol. 16</u> <u>(3): 317-20.</u> Moreau, J.L. <i>et al.</i> (1987) Monoclonal antibodies identify three epitope clusters on the mouse p55 subunit of the interleukin 2 receptor: relationship to the interleukin 2-binding site. <u>Eur J Immunol. 17 (7): 929-35.</u> Yaqoob, P. & Calder, P.C. (1997) Glutamine requirement of proliferating T lymphocytes. <u>Nutrition. 13 (7-8): 646-51.</u> Scotland, R.S. <i>et al.</i> (2011) Sex differences in resident immune cell phenotype underlie more efficient acute inflammatory responses in female mice. <u>Blood. 118 (22): 5918-27.</u> 7. Karali, D. <i>et al.</i> (2016) T cell regulation by <i>Phlomis lanata</i> protein extracts in mice. <u>Pharm Biol. 54 (2): 207-14.</u> Szuster-Ciesielska, A. <i>et al.</i> (2019) Immunogenic Evaluation of Ribosomal P-Protein Antigen P0, P1, and P2 and Pentameric Protein Complex P0-(P1-P2)₂ of <i>Plasmodium falciparum</i> in a Mouse Model. J Immunol Res. 2019: 9264217. Curina, G. <i>et al.</i> (2018) Evaluation of immune responses in mice and sheep inoculated with a live attenuated <i>Brucella melitensis.</i> REV1 vaccine produced in bioreactor. <u>Vet Immunol Immunopathol. 198: 44-53.</u> Arad, T. <i>et al.</i> (2021) CD200 -dependent and -independent immune-modulatory

	functions of neural stem cells. <u>Stem Cell Res. 56: 102559.</u> 11. Roca, C.P. <i>et al.</i> (2023) A cross entropy test allows quantitative statistical comparison of t-SNE and UMAP representations <u>Cell Reports Methods. 3 (1): 100390.</u>				
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.				
Guarantee	12 months from date of despatch				
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and re counterparts	lated U.S. and foreign			
Health And Safety Information	Material Safety Datasheet documentation #20471 available at <u>https://www.bio-rad-antibodies.com/SDS/MCA1260SBY575</u> 20471				
Regulatory	For research purposes only				

Related Products

Recommended Useful Reagents

MOUSE SEROBLOCK FcR (BUF041A) MOUSE SEROBLOCK FcR (BUF041B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	id.com	Email: antibody_sales_uk@bio-ra	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M413701:221125'

Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint