

Datasheet: MCA1260SBY800

## **BATCH NUMBER 100007984**

Description:	RAT ANTI MOUSE CD25:StarBright Yellow 800
Specificity:	CD25
Other names:	IL-2R ALPHA CHAIN
Format:	StarBright Yellow 800
Product Type:	Monoclonal Antibody
Clone:	PC61.5.3
Isotype:	IgG1
Quantity:	100 TESTS/0.5ml

# **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse	Mouse				
Product Form	Purified IgG conjugate	Purified IgG conjugated to StarBright Yellow 800 - liquid				
/lax Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm			
	StarBright Yellow 800	549	788			
reparation	Purified IgG prepared by affinity chromatography on Protein G from tissue of supernatant					
fer Solution	Phosphate buffered s	Phosphate buffered saline				
servative	0.09% Sodium Azide	(NaN <sub>3</sub> )				
abilisers	1% Bovine Serum Alb	1% Bovine Serum Albumin				
	0.1% Pluronic F68					
	0.1% PEG 3350					

B6.1 CTL cell line.

## External Database Links

#### **UniProt:**

P01590 Related reagents

#### **Entrez Gene:**

16184 Il2ra Related reagents

### **Synonyms**

II2r

#### **Fusion Partners**

Spleen cells from immunized OFA rats were fused with cells of the P3X63Ag8.653 mouse myeloma cell line.

#### **Specificity**

**Rat anti Mouse CD25 antibody, clone PC61.5.3** reacts with the low affinity alpha chain of the interleukin-2 receptor present on activated T and B cells in mice. Rat anti Mouse CD25 antibody, clone PC61.5.3 is reported to inhibit IL-2 binding and IL-2 dependent proliferation.

### Flow Cytometry

Use  $5\mu$ I of the suggested working dilution to label  $10^6$  cells in  $100\mu$ I. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

#### References

- 1. Ceredig, R. *et al.* (1985) Expression of interleukin-2 receptor as a differentiation marker on intrathymic stem cells. Nature 314: 98-100.
- 2. Lowenthal, J.W. *et al.* (1985) High and low affinity IL 2 receptors: analysis by IL 2 dissociation rate and reactivity with monoclonal anti-receptor antibody PC61. <u>J Immunol.</u> 135 (6): 3988-94.
- 3. Hashimoto, N. *et al.* (1986) Dissociation of interleukin 2-dependent and -independent B cell proliferation with monoclonal anti-interleukin 2 receptor antibody. <u>Eur J Immunol. 16</u> (3): 317-20.
- 4. Moreau, J.L. *et al.* (1987) Monoclonal antibodies identify three epitope clusters on the mouse p55 subunit of the interleukin 2 receptor: relationship to the interleukin 2-binding site. Eur J Immunol. 17 (7): 929-35.
- 5. Yaqoob, P. & Calder, P.C. (1997) Glutamine requirement of proliferating T lymphocytes. Nutrition. 13 (7-8): 646-51.
- 6. Scotland, R.S. *et al.* (2011) Sex differences in resident immune cell phenotype underlie more efficient acute inflammatory responses in female mice. <u>Blood. 118 (22): 5918-27.</u>
- 7. Karali, D. *et al.* (2016) T cell regulation by *Phlomis lanata* protein extracts in mice. Pharm Biol. 54 (2): 207-14.
- 8. Szuster-Ciesielska, A. *et al.* (2019) Immunogenic Evaluation of Ribosomal P-Protein Antigen P0, P1, and P2 and Pentameric Protein Complex P0-(P1-P2)<sub>2</sub> of *Plasmodium falciparum* in a Mouse Model. J Immunol Res. 2019: 9264217.
- 9. Curina, G. *et al.* (2018) Evaluation of immune responses in mice and sheep inoculated with a live attenuated *Brucella melitensis*. REV1 vaccine produced in bioreactor. <u>Vet Immunol Immunopathol</u>. 198: 44-53.
- 10. Arad, T. et al. (2021) CD200 -dependent and -independent immune-modulatory

of t-SNE and UMAP representations Cell Reports Methods. 3 (1): 100390. Storage Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Guarantee 12 months from date of despatch **Acknowledgements** This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts **Health And Safety** Material Safety Datasheet documentation #20471 available at: Information https://www.bio-rad-antibodies.com/SDS/MCA1260SBY800 20471 Regulatory For research purposes only

functions of neural stem cells. Stem Cell Res. 56: 102559.

11. Roca, C.P. et al. (2023) A cross entropy test allows quantitative statistical comparison

# **Related Products**

# **Recommended Useful Reagents**

MOUSE SEROBLOCK FcR (BUF041A)
MOUSE SEROBLOCK FcR (BUF041B)

**North & South** Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

**America** Fax: +1 919 878 3751

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M426484:240123'

### Printed on 15 May 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint