

Datasheet: MCA1263F

Description:	MOUSE ANTI HUMAN CD123:FITC
Specificity:	CD123
Other names:	INTERLEUKIN 3 RECEPTOR ALPHA
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	6H6
Isotype:	IgG1
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	0.2% Bovine Serum Albumin		
External Database Links	UniProt: P26951 Related reagents		
	Entrez Gene: 3563 IL3RA Related reagents		
Synonyms	IL3R		

Specificity **Mouse anti Human CD123 antibody, clone 6H6** recognizes the human interleukin-3 (IL-3) receptor alpha chain, also known as CD123. Mouse anti Human CD123 antibody, clone 6H6 is useful for monitoring expression of the IL-3 receptor on activated endothelial cells, monocytes and eosinophils, sorting bone marrow progenitor cells and purification of basophils.

Mouse anti Human CD123 antibody, clone 6H6 does not neutralize the activity of the IL-3 receptor alpha chain.

Flow Cytometry Use 5ul of the suggested working dilution to label 1 x 10⁶ cells in 100ul

References

1. Shimizu, Y. *et al.* (2008) Interleukin-3 does not affect the differentiation of mast cells derived from human bone marrow progenitors. [Immunol Invest.37: 1-17](#)
2. Massone, C. *et al.* (2011) Immunophenotype of skin lymphocytic infiltrate in patients co-infected with Mycobacterium leprae and human immunodeficiency virus: a scenario dependent on CD8+ and/or CD20+ cells. [Br J Dermatol. 165: 321-8.](#)

Further Reading

1. Korpelainen, E.I. *et al.* (1993) The receptor for interleukin 3 is selectively induced in human endothelial cells by tumor necrosis factor alpha and potentiates interleukin 8 secretion and neutrophil transmigration. [Proc Natl Acad Sci U S A. 90 \(23\): 11137-41.](#)

Storage Store at +4°C. DO NOT FREEZE.
This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M344241:190111'

Printed on 11 Aug 2020