

# Datasheet: MCA1266 BATCH NUMBER 1711

Description:	MOUSE ANTI MOUSE CD161 / NK1.1
Specificity:	CD161 / NK1.1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	PK136
lsotype:	lgG2a
Quantity:	0.25 mg

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal											
	communications from the	communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>										
	information. For general											
	rad-antibodies.com/protocols.											
		Yes	No	Not Determined	Suggested Dilution							
	Flow Cytometry	-			5ug/ml - 10ug/ml							
	Immunohistology - Frozen			•								
	Immunohistology - Paraffin											
	ELISA			•								
	Immunoprecipitation	-										
	Western Blotting											
	Cytotoxic Assays	-			1ug/ml - 10ug/ml							
	Where this antibody has not been tested for use in a particular technique this does not											
	necessarily exclude its us	necessarily exclude its use in such procedures. Suggested working dilutions are given as										
	•	a guide only. It is recommended that the user titrates the antibody for use in their own										
	system using appropriate negative/positive controls. Clone PK136 has also been reported to be suitable for <i>in vivo</i> depletion of NK cells.											
							Target Species	Mouse				
							Species Cross Reactivity	Does not react with:Rat, Human				
Product Form	Purified IgG - liquid Purified IgG prepared by affinity chromatography on Protein A											
Preparation												
Buffer Solution	Phosphate buffered saline											

Preservative Stabilisers	0.09% Sodium Azide		
Carrier Free	Yes		
Approx. Protein Concentrations	IgG concentration 1 mg/ml		
Immunogen	Spleen and bone marrow cells from CE mice.		
External Database Links	UniProt:P27814Related reagentsP27812Related reagentsEntrez Gene:17059Klrb1cRelated reagents80782Klrb1bRelated reagents		
Synonyms	Ly55b, Ly55c, Nkrp1b, Nkrp1c		
RRID	AB_322347		
Fusion Partners	Spleen cells from immunised (C3H x BALB/c) FI Hybrid were fused with cells of the Sp2/0 - Ag14 myeloma cell line.		
Specificity	<ul> <li>Mouse anti Mouse CD161 / NK1.1 antibody, clone PK136 recognizes the mouse NK1.1 cell surface antigen, a cell surface glycoprotein encoded by members of the NKR-P1 gene family. The NK1.1 surface antigen is also known as CD161b/CD161c and Ly-55.</li> <li>In the mouse the NKR-P1 family has three members, NKR-P1A, -B and -C, whilst in the human only one member has been identified. The human protein has received the designation CD161, and the mouse proteins have been referred to as CD161a, -b, -c etc.</li> <li>Although previously thought to recognize only CD161c, recent data has shown that the PK136 antibody may also react with CD161b. CD161c expression itself is strain specific in mice, but recognition of CD161b by PK136 appears to be even more complex, as only some CD161b positive strains are labelled by the antibody. Engagement of CD161c has been reported to have activating function in NK cells, whilst engagement of CD161b is inhibitory.</li> <li>Mouse anti Mouse NK1.1 Antigen antibody, clone PK136 is useful for the identification of NK cells in selected strains of mice (positive on C57BL, FVB/N and NZB, but negative on AKR and BALB/c) and is also expressed by rare subsets of T cells and monocytes. Mouse anti Mouse NK1.1 antibody, clone PK136 has also been used for <i>in vivo</i> depletion of NK</li> </ul>		
Flow Cytometry	cells and <i>in vitro</i> activation of NK cells. Use 10ul of the suggested working dilution to label 1 x 10 <sup>6</sup> cells in 100ul.		

1. Wang, M. *et al.* (1998) Natural killer cell depletion fails to influence initial CD4 T cell commitment in vivo in exogenous antigen-stimulated cytokine and antibody responses. J Immunol. 160 (3): 1098-105.

2. Koo, G.C. *et al.* (1986) The NK-1.1(-) mouse: a model to study differentiation of murine NK cells. <u>J Immunol. 137 (12): 3742-7.</u>

3. Kung, S.K. *et al.* (1999) The NKR-P1B gene product is an inhibitory receptor on SJL/J NK cells. J Immunol. 162 (10): 5876-87.

4. Carlyle, J.R. *et al.* (1999) Mouse NKR-P1B, a novel NK1.1 antigen with inhibitory function. J Immunol. 162 (10): 5917-23.

5. Carnemolla, B. *et al.* (2002) Enhancement of the antitumor properties of interleukin-2 by its targeted delivery to the tumor blood vessel extracellular matrix. <u>Blood. 99: 1659-65.</u>

6. Carpentier, A.F. *et al.* (1999) Oligodeoxynucleotides containing CpG motifs can induce rejection of a neuroblastoma in mice. <u>Cancer Res. 59: 5429-32.</u>

7. Sakai, T. *et al.* (2010) Inflammatory disease and cancer with a decrease in Kupffer cell numbers in Nucling-knockout mice. Int J Cancer. 126: 1079-94.

8. Svensson, L. *et al.* (2003) gammadelta T cells contribute to the systemic immunoglobulin E response and local B-cell reactivity in allergic eosinophilic airway inflammation. Immunology. 108 (1): 98-108.

9. Hazlett, L.D. *et al.* (2007) NKT cells are critical to initiate an inflammatory response after *Pseudomonas aeruginosa* ocular infection in susceptible mice. <u>J Immunol. 179 : 1138-46.</u>

10. Joseph-Pietras, D. *et al.* (2006) Anti-tumoural activity of peripheral blood mononuclear cells against melanoma cells: discrepant in-vitro and in-vivo effects. <u>Melanoma Res. 16:</u> <u>325-33.</u>

11. Gock, H. *et al.* (2014) Altered glycosylation in donor mice causes rejection of strainmatched skin and heart grafts. <u>Am J Transplant. 14 (4): 797-805.</u>

12. Khallouf, H. *et al.* (2012) 5-Fluorouracil and interferon- $\alpha$  immunochemotherapy enhances immunogenicity of murine pancreatic cancer through upregulation of NKG2D ligands and MHC class I. J Immunother. 35 (3): 245-53.

13. Ebbinghaus, C. *et al.* (2005) Engineered vascular-targeting antibody-interferon-gamma fusion protein for cancer therapy. <u>Int J Cancer. 116 (2): 304-13.</u>

14. Ekstrand-Hammarström, B. *et al.* (2011) Inhalation of alkylating mustard causes long-term T cell-dependent inflammation in airways and growth of connective tissue. <u>Toxicology. 280 (3): 88-97.</u>

15. Klezovich-Bénard M *et al.* (2012) Mechanisms of NK cell-macrophage *Bacillus anthracis* crosstalk: a balance between stimulation by spores and differential disruption by toxins. <u>PLoS Pathog. 8 (1): e1002481.</u>

16. Halin, C. *et al.* (2002) Enhancement of the antitumor activity of interleukin-12 by targeted delivery to neovasculature. <u>Nat Biotechnol. 20 (3): 264-9.</u>

17. Gustafsson, Å. *et al.* (2015) Differential cellular responses in healthy mice and in mice with established airway inflammation when exposed to hematite nanoparticles. <u>Toxicol</u> <u>Appl Pharmacol. 288 (1): 1-11.</u>

18. Flavell, D.J. *et al.* (2019) The TLR3 Agonist Poly Inosinic:Cytidylic Acid Significantly Augments the Therapeutic Activity of an Anti-CD7 Immunotoxin for Human T-cell Leukaemia. <u>Biomedicines. 7 (1) Feb 16 [Epub ahead of print].</u>

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1266 10040
Regulatory	For research purposes only

### **Related Products**

### **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87	) <u>HRP</u>
Goat Anti Mouse IgG (STAR76)	RPE
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>
Goat Anti Mouse IgG (H/L) (STAR117)	<u>Alk. Phos.</u> , <u>DyLight®488</u> , <u>DyLight®550</u> ,
	<u>DyLight®650</u> , <u>DyLight®680</u> , <u>DyLight®800</u> ,
	FITC, HRP
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>
Goat Anti Mouse IgG (STAR77)	HRP
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP
Rabbit Anti Mouse IgG (STAR13)	HRP
 arth & Cauth Tal. 1 900 265 7276	

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M365022:200529'

#### Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint