

Datasheet: MCA1266A647

## **BATCH NUMBER 1707**

Description:	MOUSE ANTI MOUSE CD161 / NK1.1:Alexa Fluor® 647
Specificity:	CD161 / NK1.1
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	PK136
Isotype:	IgG2a
Quantity:	100 TESTS/1ml

# **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse					
Species Cross Reactivity	Does not react with:	Does not react with:Rat, Human				
Product Form	Purified IgG conjugated to Alexa Fluor 647 - liquid					
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm			
	Alexa Fluor®647	650	665			
Preparation	Purified IgG prepare	Purified IgG prepared by affinity chromatography on Protein A				
Buffer Solution	Phosphate buffered	saline				
Preservative	0.09% Sodium Azide (NaN <sub>3</sub> )					
Stabilisers	1% Bovine Serum Albumin					
Approx. Protein	IgG concentration 0.	05 mg/ml				

Concentrations	
Immunogen	Spleen and bone marrow cells from CE mice.
External Database Links	UniProt: P27814 Related reagents P27812 Related reagents
	Entrez Gene:
	17059Klrb1cRelated reagents80782Klrb1bRelated reagents
Synonyms	Ly55b, Ly55c, Nkrp1b, Nkrp1c
Fusion Partners	Spleen cells from immunised (C3H x BALB/c) FI Hybrid were fused with cells of the Sp2/0 - Ag14 myeloma cell line.
Specificity	Mouse anti Mouse CD161 / NK1.1 antibody, clone PK136 recognizes the mouse NK1.1 cell surface antigen, a cell surface glycoprotein encoded by members of the NKR-P1 gene family. The NK1.1 surface antigen is also known as CD161b/CD161c and Ly-55.  In the mouse the NKR-P1 family has three members, NKR-P1A, -B and -C, whilst in the human only one member has been identified. The human protein has received the designation CD161, and the mouse proteins have been referred to as CD161a, -b, -c etc.
	Although previously thought to recognize only CD161c, recent data has shown that the PK136 antibody may also react with CD161b. CD161c expression itself is strain specific in mice, but recognition of CD161b by PK136 appears to be even more complex, as only some CD161b positive strains are labelled by the antibody. Engagement of CD161c has been reported to have activating function in NK cells, whilst engagement of CD161b is inhibitory.
	Mouse anti Mouse NK1.1 Antigen antibody, clone PK136 is useful for the identification of NK cells in selected strains of mice (positive on C57BL, FVB/N and NZB, but negative on AKR and BALB/c) and is also expressed by rare subsets of T cells and monocytes. Mouse anti Mouse NK1.1 antibody, clone PK136 has also been used for <i>in vivo</i> depletion of NK cells and <i>in vitro</i> activation of NK cells.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
	The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR ( <u>BUF041A/B</u> )
References	1 Wang M et al. (1998) Natural killer cell depletion fails to influence initial CD4 T cell

# References

- 1. Wang, M. et al. (1998) Natural killer cell depletion fails to influence initial CD4 T cell commitment in vivo in exogenous antigen-stimulated cytokine and antibody responses.  $\underline{\mathsf{J}}$ Immunol. 160 (3): 1098-105.
- 2. Koo, G.C. et al. (1986) The NK-1.1(-) mouse: a model to study differentiation of murine

NK cells. J Immunol. 137 (12): 3742-7.

- 3. Kung, S.K. *et al.* (1999) The NKR-P1B gene product is an inhibitory receptor on SJL/J NK cells. J Immunol. 162 (10): 5876-87.
- 4. Carlyle, J.R. *et al.* (1999) Mouse NKR-P1B, a novel NK1.1 antigen with inhibitory function. <u>J Immunol</u>. 162 (10): 5917-23.
- 5. Carnemolla, B. *et al.* (2002) Enhancement of the antitumor properties of interleukin-2 by its targeted delivery to the tumor blood vessel extracellular matrix. <u>Blood. 99: 1659-65.</u>
- 6. Carpentier, A.F. *et al.* (1999) Oligodeoxynucleotides containing CpG motifs can induce rejection of a neuroblastoma in mice. <u>Cancer Res. 59: 5429-32.</u>
- 7. Sakai, T. *et al.* (2010) Inflammatory disease and cancer with a decrease in Kupffer cell numbers in Nucling-knockout mice. Int J Cancer. 126: 1079-94.
- 8. Svensson, L. *et al.* (2003) gammadelta T cells contribute to the systemic immunoglobulin E response and local B-cell reactivity in allergic eosinophilic airway inflammation. Immunology. 108 (1): 98-108.
- 9. Hazlett, L.D. *et al.* (2007) NKT cells are critical to initiate an inflammatory response after *Pseudomonas aeruginosa* ocular infection in susceptible mice. <u>J Immunol. 179:</u> 1138-46.
- 10. Joseph-Pietras, D. *et al.* (2006) Anti-tumoural activity of peripheral blood mononuclear cells against melanoma cells: discrepant in-vitro and in-vivo effects. <u>Melanoma Res. 16:</u> 325-33.
- 11. Gock, H. *et al.* (2014) Altered glycosylation in donor mice causes rejection of strain-matched skin and heart grafts. Am J Transplant. 14 (4): 797-805.
- 12. Khallouf, H. *et al.* (2012) 5-Fluorouracil and interferon-α immunochemotherapy enhances immunogenicity of murine pancreatic cancer through upregulation of NKG2D ligands and MHC class I. <u>J Immunother. 35 (3): 245-53.</u>
- 13. Ebbinghaus, C. *et al.* (2005) Engineered vascular-targeting antibody-interferon-gamma fusion protein for cancer therapy. <u>Int J Cancer. 116 (2): 304-13.</u>
- 14. Ekstrand-Hammarström, B. *et al.* (2011) Inhalation of alkylating mustard causes long-term T cell-dependent inflammation in airways and growth of connective tissue. <u>Toxicology. 280 (3): 88-97.</u>
- 15. Klezovich-Bénard M *et al.* (2012) Mechanisms of NK cell-macrophage *Bacillus anthracis* crosstalk: a balance between stimulation by spores and differential disruption by toxins. <u>PLoS Pathog. 8 (1): e1002481.</u>
- 16. Halin, C. *et al.* (2002) Enhancement of the antitumor activity of interleukin-12 by targeted delivery to neovasculature. <u>Nat Biotechnol. 20 (3): 264-9.</u>
- 17. Gustafsson, Å. *et al.* (2015) Differential cellular responses in healthy mice and in mice with established airway inflammation when exposed to hematite nanoparticles. <u>Toxicol Appl Pharmacol. 288 (1): 1-11.</u>
- 18. Flavell, D.J. *et al.* (2019) The TLR3 Agonist Poly Inosinic:Cytidylic Acid Significantly Augments the Therapeutic Activity of an Anti-CD7 Immunotoxin for Human T-cell Leukaemia. <u>Biomedicines</u>. 7 (1) Feb 16 [Epub ahead of print].

### **Storage**

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this

Guarantee	12 months from date of despatch	
Acknowledgements	This product is provided under an intellectual property licence of Corporation. The transfer of this product is contingent on the beginned product solely in research, excluding contract research or any stand the buyer must not sell or otherwise transfer this product of diagnostic, therapeutic or prophylactic purposes; (b) testing, and services, or information in return for compensation on a per-test or quality assurance or quality control, or (d) resale, whether or research. For information on purchasing a license to this product as described above, contact Life Technologies Corporation, 57 CA 92008 USA or outlicensing@thermofisher.com	uyer using the purchased fee for service research, or its components for (a) halysis or screening st basis; (c) manufacturing or not resold for use in act for purposes other than
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1266A647">https://www.bio-rad-antibodies.com/SDS/MCA1266A647</a> 10041	
Regulatory	For research purposes only	

# **Related Products**

# **Recommended Useful Reagents**

MOUSE SEROBLOCK FcR (BUF041A)
MOUSE SEROBLOCK FcR (BUF041B)

North & South Tel: +1 800 265 7376

America

Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M365023:200529'

# Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint