

Datasheet: MCA1267P647T

Description:	MOUSE ANTI HUMAN CD4:RPE-Alexa Fluor® 647
Specificity:	CD4
Format:	RPE-ALEXA FLUOR® 647
<b>Product Type:</b>	Monoclonal Antibody
Clone:	RPA-T4
Isotype:	lgG1
Quantity:	25 TESTS

# **Product Details**

## **Applications**

Links

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="https://www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human						
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - Alexa Fluor® 647 - lyophilized						
Reconstitution	Reconstitute in 0.25 ml disilled water						
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)				
	RPE-Alexa Fluor®647 488nm laser	496	667				
	RPE-Alexa Fluor®647 561nm laser	546	667				
Preparation	Purified IgG prepared	by affinity chromatog	raphy on Protein G fro	om tissue culture supernatant			
Buffer Solution	Phosphate buffered s	aline					
Preservative	0.09% Sodium Azide						
Stabilisers	1% Bovine Serum Albumin						
	5% Sucrose						
Immunogen	Human PHA blasts						
External Database	Ha:Duct.						

**UniProt:** 

P01730 Related reagents

#### **Entrez Gene:**

920 CD4 Related reagents

#### **Fusion Partners**

Spleen cells from immunised BALB/c mice were fused with cells of the mouse NSI myeloma cell line

## **Specificity**

**Mouse anti human CD4 antibody, clone RPA-T4** recognizes human CD4, a 55 kDa cell surface glycoprotein that is primarily expressed on a subpopulation of T lymphocytes, on peripheral blood monocytes and on tissue macrophages. Epitope mapping studies have shown that antibodies, produced by clone RPA-T4, recognize an epitope within domain 1 of the extracellular region of the CD4 molecule.

Mouse anti human CD4 antibody, clone RPA-T4 has been reported to block gp120-CD4 interaction and inhibit syncytium formation, see Piatier-Tonneau *et al*, 1997 for details. Bio-Rad recommend the use of Mouse anti Human CD4:Low Endotoxin (MCA1267EL) for this purpose.

# **Flow Cytometry**

Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells or 100ul whole blood. Tandem conjugates of RPE-Alexa Fluor® 647 display fluorescence properties that are virtually identical to those of RPE-Cy5 tandem conjugates.

#### References

- 1. Zarkesh-Esfahani, H. *et al.* (2001) High-dose leptin activates human leukocytes via receptor expression on monocytes. J Immunol. 167 (8): 4593-9.
- 2. Voehringer, D. *et al.* (2002) Lack of proliferative capacity of human effector and memory T cells expressing killer cell lectinlike receptor G1 (KLRG1). <u>Blood. 100 (10): 3698-702.</u>
- 3. Piatier-Tonneau, D. (1997) CD4 workshop panel report. In: Leucocyte Typing VI: White Cell Differentiation Antigens: Proceedings of the Sixth International Workshop and Conference Held in Kobe, Japan, 10-14 November 1996. Garland Pub., 1998.
- 4. Pentón-Rol, G. et al. (2011) C-Phycocyanin ameliorates experimental autoimmune encephalomyelitis and induces regulatory T cells. Int Immunopharmacol. 11 (1): 29-38.
- 5. Wright, G.J. *et al.* (2001) The unusual distribution of the neuronal/lymphoid cell surface CD200 (OX2) glycoprotein is conserved in humans. <u>Immunology</u>. 102 (2): 173-9.
- 6. Zhang, Y. et al. (2013) Accelerated *In Vivo* Proliferation of Memory Phenotype CD4+ T-cells in Human HIV-1 Infection Irrespective of Viral Chemokine Co-receptor PLoS Pathog 9(4): e1003310.
- 7. Bughani, U. *et al.* (2017) T cell activation and differentiation is modulated by a CD6 domain 1 antibody Itolizumab. PLoS One. 12 (7): e0180088.

## Storage

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

## DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

### Shelf Life

12 months from date of reconstitution.

## **Acknowledgements**

This product is provided under an intellectual property license from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchased product solely in research conducted by the buyer, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in

return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com

Health And Safety
Information

Material Safety Datasheet documentation #10075 available at:

10075: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf</a>

**Regulatory** For research purposes only

# Related Products

# **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL:RPE-Alexa Fluor® 647 (MCA928P647)

## **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376 Worldwide
America Fax: +1 919 878 3751

Email: antibody\_sales\_us@bio-rad.com

Tel: +44 (0)1865 852 700 **Europe** Fax: +44 (0)1865 852 739

Email: antibody\_sales\_uk@bio-rad.com

'M322590:180727'

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_de@bio-rad.com

Printed on 06 Dec 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint