

# Datasheet: MCA1267SBB700T

**BATCH NUMBER 100007474**

<b>Description:</b>	MOUSE ANTI HUMAN CD4:StarBright Blue 700
<b>Specificity:</b>	CD4
<b>Format:</b>	StarBright Blue 700
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	RPA-T4
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	25 TESTS/0.125ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human		
<b>Product Form</b>	Purified IgG conjugated to StarBright Blue 700 - liquid		
<b>Max Ex/Em</b>	<b>Fluorophore</b>	<b>Excitation Max (nm)</b>	<b>Emission Max (nm)</b>
	StarBright Blue 700	470	700
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
<b>Buffer Solution</b>	Phosphate buffered saline		
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )		
	1% Bovine Serum Albumin		
	0.1% Pluronic F68		
	0.1% PEG 3350		
	0.05% Tween 20		

<b>Immunogen</b>	Human PHA blasts
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P01730</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">920</a>    CD4    <a href="#">Related reagents</a></p>
<b>Fusion Partners</b>	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NSI myeloma cell line
<b>Specificity</b>	<p><b>Mouse anti human CD4 antibody, clone RPA-T4</b> recognizes human CD4, a ~55 kDa cell surface glycoprotein, primarily expressed on a subpopulation of T lymphocytes, on peripheral blood monocytes and on tissue macrophages. Epitope mapping shows that antibodies, produced by clone RPA-T4, recognize an epitope within domain 1 of the extracellular region of the CD4 molecule.</p> <p>Mouse anti human CD4 antibody, clone RPA-T4 blocks gp120-CD4 interaction and inhibits syncytium formation (Piatier-Tonneau <i>et al</i>, 1997). The use of <a href="#">Mouse anti Human CD4:Low Endotoxin (MCA1267EL)</a> is recommended for functional assays.</p>
<b>Flow Cytometry</b>	Use 5µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Zarkesh-Esfahani, H. <i>et al.</i> (2001) High-dose leptin activates human leukocytes via receptor expression on monocytes. <a href="#">J Immunol. 167 (8): 4593-9.</a></li> <li>2. Voehringer, D. <i>et al.</i> (2002) Lack of proliferative capacity of human effector and memory T cells expressing killer cell lectinlike receptor G1 (KLRG1). <a href="#">Blood. 100 (10): 3698-702.</a></li> <li>3. Piatier-Tonneau, D. (1997) CD4 workshop panel report. In: Leucocyte Typing VI: White Cell Differentiation Antigens: Proceedings of the Sixth International Workshop and Conference Held in Kobe, Japan, 10-14 November 1996. Garland Pub., 1998.</li> <li>4. Pentón-Rol, G. <i>et al.</i> (2011) C-Phycocyanin ameliorates experimental autoimmune encephalomyelitis and induces regulatory T cells. <a href="#">Int Immunopharmacol. 11 (1): 29-38.</a></li> <li>5. Wright, G.J. <i>et al.</i> (2001) The unusual distribution of the neuronal/lymphoid cell surface CD200 (OX2) glycoprotein is conserved in humans. <a href="#">Immunology. 102 (2): 173-9.</a></li> <li>6. Zhang, Y. <i>et al.</i> (2013) Accelerated <i>in vivo</i>. proliferation of memory phenotype CD4<sup>+</sup> T-cells in human HIV-1 infection irrespective of viral chemokine co-receptor tropism. <a href="#">PLoS Pathog. 9 (4): e1003310.</a></li> <li>7. Bughani, U. <i>et al.</i> (2017) T cell activation and differentiation is modulated by a CD6 domain 1 antibody Itolizumab. <a href="#">PLoS One. 12 (7): e0180088.</a></li> </ol>
<b>Storage</b>	<p>Store at +4°C. DO NOT FREEZE.</p> <p>This product should be stored undiluted.</p>
<b>Guarantee</b>	12 months from date of despatch
<b>Acknowledgements</b>	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign

counterparts

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #20471 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1267SBB700T">https://www.bio-rad-antibodies.com/SDS/MCA1267SBB700T</a> 20471
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M415443:221223'

**Printed on 18 Apr 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)