Datasheet: MCA1267SBUV605T

BATCH NUMBER 64718177

Description:	MOUSE ANTI HUMAN CD4:StarBright UltraViolet 605
Specificity:	CD4
Format:	StarBright UltraViolet 605
Product Type:	Monoclonal Antibody
Clone:	RPA-T4
Isotype:	IgG1
Quantity:	25 TESTS/0.125ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Human

Product Form

Purified IgG conjugated to StarBright UltraViolet 605 - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
StarBright UltraViolet 605	340	609

Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

0.09% Sodium Azide (NaN_3)
1% Bovine Serum Albumin
0.1% Pluronic F68
0.1% PEG 3350

	0.05% Tween 20
Approx. Protein Concentrations	For information on the concentration of our StarBright Dye conjugated reagents please visit our FAQ page.
Immunogen	Human PHA blasts
External Database Links	**UniProt:** P01730 Related reagents **Entrez Gene:** 920 CD4 Related reagents
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NSI myeloma cell line
Specificity	**Mouse anti human CD4 antibody, clone RPA-T4** recognizes human CD4, a ~55 kDa cell surface glycoprotein, primarily expressed on a subpopulation of T lymphocytes, on peripheral blood monocytes and on tissue macrophages. Epitope mapping shows that antibodies, produced by clone RPA-T4, recognize an epitope within domain 1 of the extracellular region of the CD4 molecule. Mouse anti human CD4 antibody, clone RPA-T4 blocks gp120-CD4 interaction and inhibits syncytium formation (Piatier-Tonneau *et al*, 1997).
Flow Cytometry	Use 5µl of the suggested working dilution to label 10^6 cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
References	1. Piatier-Tonneau, D. (1997) CD4 workshop panel report. In: Leucocyte Typing VI: White Cell Differentiation Antigens: Proceedings of the Sixth International Workshop and Conference Held in Kobe, Japan, 10-14 November 1996. Garland Pub., 1998. 2. Zarkesh-Esfahani, H. *et al.* (2001) High-dose leptin activates human leukocytes via receptor expression on monocytes. J Immunol. 167 (8): 4593-9. 3. Wright, G.J. *et al.* (2001) The unusual distribution of the neuronal/lymphoid cell surface CD200 (OX2) glycoprotein is conserved in humans. Immunology. 102 (2): 173-9. 4. Pentón-Rol, G. *et al.* (2011) C-Phycocyanin ameliorates experimental autoimmune encephalomyelitis and induces regulatory T cells. Int Immunopharmacol. 11 (1): 29-38. 5. Zhang, Y. *et al.* (2013) Accelerated *in vivo.* proliferation of memory phenotype $CD4^+$ T-cells in human HIV-1 infection irrespective of viral chemokine co-receptor tropism. PLoS Pathog. 9 (4): e1003310. 6. Bughani, U. *et al.* (2017) T cell activation and differentiation is modulated by a CD6 domain 1 antibody Itolizumab. PLoS One. 12 (7): e0180088. 7. Agrawal, S.M. *et al.* (2013) Extracellular matrix metalloproteinase inducer shows active perivascular cuffs in multiple sclerosis. Brain. 136 (Pt 6): 1760-77. 8. Malmassari, S.L. *et al.* (2007) Impact of hepatitis B virus basic core promoter mutations on T cell response to an immunodominant HBx-derived epitope. Hepatology. 45 (5): 1199-209.

9. Wooldridge, L. *et al.* (2006) Anti-coreceptor antibodies profoundly affect staining with peptide-MHC class I and class II tetramers. Eur J Immunol. 36 (7): 1847-55.
10. Wildum, S. *et al.* (2006) Contribution of Vpu, Env, and Nef to CD4 down-modulation and resistance of human immunodeficiency virus type 1-infected T cells to superinfection. J Virol. 80 (16): 8047-59.
11. Kirchhof, J. *et al.* (2018) Learned immunosuppressive placebo responses in renal transplant patients. Proc Natl Acad Sci U S A. 115 (16): 4223-7.
12. Kelleher, M. *et al.* (2011) Comparative Kinetics of Immune Responses and Changes in Cellular Sub-Sets Detected in Colorectal Cancer Patients Vaccinated with MVA-5T4 (TroVax) Administered Alongside Two Different Chemotherapy Regimens J Cancer Therapy. 02 (01): 54-64.
13. Turuntaš, V. *et al.* (2024) The Effect of Static Magnetic Fields of Different Strengths and Polarities on Cytokine Production by Human Lymphocytes *In Vitro* Bioengineering. 11 (8): 749.

Storage	This product is shipped at ambient temperature. Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA1267SBUV605T
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A)
HUMAN SEROBLOCK (BUF070B)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M439818:250523'

Printed on 28 May 2026