

# Datasheet: MCA1267SBV475T BATCH NUMBER 64596747

Description:	MOUSE ANTI HUMAN CD4:StarBright Violet 475
Specificity:	CD4
Format:	StarBright Violet 475
Product Type:	Monoclonal Antibody
Clone:	RPA-T4
Isotype:	lgG1
Quantity:	25 TESTS/0.125ml

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry	-			Neat	
	Where this product has necessarily exclude its a guide only. It is recon system using appropria	use in such pr nmended that	ocedur	res. Suggested workir er titrates the product	ng dilutions are given as	
Target Species	Human					
Product Form	Purified IgG conjugated to StarBright Violet 475 - liquid					
Max Ex/Em	Fluorophore StarBright Violet 475	Excitation Max 405	(nm)	Emission Max (nm) 479		
Preparation	Purified IgG prepared b supernatant	by affinity chroi	matogr	aphy on Protein G fro	m tissue culture	
Buffer Solution	Phosphate buffered saline					
Preservative	0.09% Sodium Azide (NaN <sub>3</sub> )					
Stabilisers	1% Bovine Serum Albumin					
	0.1% Pluronic F68					
	0.1% PEG 3350					
	0.05% Tween 20					

Immunogen	Human PHA blasts				
External Database Links	UniProt:         P01730       Related reagents         Entrez Gene:         920       CD4         Related reagents				
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NSI myeloma cell line				
Specificity	Mouse anti human CD4 antibody, clone RPA-T4 recognizes human CD4, a ~55 kDa cell surface glycoprotein, primarily expressed on a subpopulation of T lymphocytes, on peripheral blood monocytes and on tissue macrophages. Epitope mapping shows that antibodies, produced by clone RPA-T4, recognize an epitope within domain 1 of the extracellular region of the CD4 molecule.				
Flow Cytometry	syncytium formation ( <u>Piatier-Tonneau <i>et al</i>, 1997</u> ). Use 5µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.				
References	<ol> <li>Piatier-Tonneau, D. (1997) CD4 workshop panel report. In: Leucocyte Typing VI: White Cell Differentiation Antigens: Proceedings of the Sixth International Workshop and Conference Held in Kobe, Japan, 10-14 November 1996. Garland Pub., 1998.</li> <li>Zarkesh-Esfahani, H. <i>et al.</i> (2001) High-dose leptin activates human leukocytes via receptor expression on monocytes. J Immunol. 167 (8): 4593-9.</li> <li>Wright, G.J. <i>et al.</i> (2001) The unusual distribution of the neuronal/lymphoid cell surface CD200 (OX2) glycoprotein is conserved in humans. Immunology. 102 (2): 173-9.</li> <li>Pentón-Rol, G. <i>et al.</i> (2011) C-Phycocyanin ameliorates experimental autoimmune encephalomyelitis and induces regulatory T cells. Int Immunopharmacol. 11 (1): 29-38.</li> <li>Zhang, Y. <i>et al.</i> (2013) Accelerated <i>in vivo</i>. proliferation of memory phenotype CD4<sup>+</sup> T-cells in human HIV-1 infection irrespective of viral chemokine co-receptor tropism. PLoS Pathog. 9 (4): e1003310.</li> <li>Bughani, U. <i>et al.</i> (2017) T cell activation and differentiation is modulated by a CD6 domain 1 antibody Itolizumab. PLoS One. 12 (7): e0180088.</li> <li>Agrawal, S.M. <i>et al.</i> (2007) Impact of hepatitis B virus basic core promoter mutations on T cell response to an immunodominant HBx-derived epitope. Hepatology. 45 (5): 1199-209.</li> <li>Wooldridge, L. <i>et al.</i> (2006) Anti-coreceptor antibodies profoundly affect staining with peptide-MHC class I and class II tetramers. Eur J Immunol. 36 (7): 1847-55.</li> <li>Wildum, S. <i>et al.</i> (2006) Contribution of Vpu, Env, and Nef to CD4 down-modulation and resistance of human immunodeficiency virus type 1-infected T cells to superinfection. J Virol. 80 (16): 8047-59.</li> </ol>				

	<ul> <li>11. Kirchhof, J. <i>et al.</i> (2018) Learned immunosuppressive placebo responses in renal transplant patients. <u>Proc Natl Acad Sci U S A. 115 (16): 4223-7.</u></li> <li>12. Kelleher, M. <i>et al.</i> (2011) Comparative Kinetics of Immune Responses and Changes in Cellular Sub-Sets Detected in Colorectal Cancer Patients Vaccinated with MVA-5T4 (TroVax) Administered Alongside Two Different Chemotherapy Regimens <u>J Cancer Therapy. 02 (01): 54-64.</u></li> </ul>			
Storage	Store at +4°C. DO NOT FREEZE.			
	This product should be stored undiluted.			
Guarantee	12 months from date of despatch			
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and rel counterparts	ated U.S. and foreign		
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA1267SBV475T 20471			
Regulatory	For research purposes only			

### **Related Products**

#### **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M415455:221223'

#### Printed on 27 Jun 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint