

## Datasheet: MCA1269F

<b>Description:</b>	MOUSE ANTI HUMAN CD100:FITC
<b>Specificity:</b>	CD100
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	A8
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

### Target Species

Human

### Species Cross Reactivity

Reacts with: Cynomolgus monkey, Rhesus Monkey

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

### Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid from tissue culture supernatant.

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

### Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

### Buffer Solution

Phosphate buffered saline

<b>Preservative</b>	0.09% Sodium Azide
<b>Stabilisers</b>	1% Bovine Serum Albumin
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1 mg/ml
<b>Immunogen</b>	Human PHA activated lymphocytes.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">Q92854</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">10507</a> SEMA4D    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	C9orf164, CD100, SEMAJ
<b>RRID</b>	AB_324619
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
<b>Specificity</b>	<b>Mouse anti Human CD100 antibody, clone A8</b> recognizes human Sepaphorin 4D, also known as CD100, A8, BB18 or GR3. CD100 is a 841 amino acid ~150 kDa homodimeric single pass type 1 transmembrane cell surface glycoprotein acting as a receptor for some members of the plexin family. CD100 is expressed at low levels by peripheral blood T cells, and at higher density on activated T cells and B cells, it is also expressed at high levels in fetal kidney with lower levels in fetal brain, liver and lung ( <a href="#">Maestrini et al. 1996</a> ).
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Aversa, G.G. <i>et al.</i> (1989) Use of monoclonal antibodies to study <i>in vivo</i> and <i>in vitro</i>-activated lymphocytes. <a href="#">Transplant Proc. 21 (1 Pt 1): 349-50.</a></li> <li>2. Van Vliet, S. J. <i>et al.</i> (1995) Species cross reactivity (human-monkey-bovine) of the adhesion structure section mAb. Leucocyte Typing V. Oxford University Press p1607-8.</li> <li>3. Mizrahi, S. <i>et al.</i> (2007) CD100 on NK cells enhance IFN<math>\gamma</math> secretion and killing of target cells expressing CD72. <a href="#">PLoS One. 2 (9): e818.</a></li> <li>4. Yoshino, N. <i>et al.</i> (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (<i>Macaca fascicularis</i>) by using anti-human cross-reactive antibodies. <a href="#">Exp Anim. 49 (2): 97-110.</a></li> </ol>
<b>Storage</b>	<p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.</p>

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10041 available at: 10041: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf</a>
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

From March 15, 2021, we will no longer supply printed datasheets with our products.  
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](https://www.bio-rad-antibodies.com)

'M374411:201103'

Printed on 09 Feb 2021