

Datasheet: MCA1270

BATCH NUMBER 166543

Description:	MOUSE ANTI HUMAN CD13
Specificity:	CD13
Other names:	AMINOPEPTIDASE N
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	WM15
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation	▪			
Western Blotting		▪		

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	<p>Reacts with: Rhesus Monkey</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture

supernatant

Buffer Solution Phosphate buffered saline

Preservative Stabilisers 0.09% sodium azide (NaN₃)

Carrier Free Yes

Approx. Protein Concentrations IgG concentration 1.0 mg/ml

Immunogen Human AML cells.

External Database Links

UniProt:
[P15144](#) [Related reagents](#)

Entrez Gene:
[290](#) ANPEP [Related reagents](#)

Synonyms APN, CD13, PEPN

RRID AB_321309

Fusion Partners Spleen cells from immunized BALB/c mice where fused with cells of the mouse NS1 myeloma cell line.

Specificity **Mouse anti Human CD13 antibody, clone WM15** recognizes human CD13 also known as aminopeptidase N. CD13 is a single pass type II glycosylated integral membrane protein with a predicted molecular mass of ~110 kDa and an apparent molecular mass of ~150 kDa expressed by granulocytes, monocytes, fibroblasts, endothelial cells and by myeloid leukaemia cells ([Bradstock *et al.* 1985](#)). CD13 acts as a major cell surface receptor for group 1 coronaviruses ([Breslin *et al.* 2003](#)) which bind to a critical sequence encompassing amino acid residues 288-295 ([Kolb *et al.* 1997](#)).

CD13 functions as an [aminopeptidase](#) enzyme, a metalloprotease present as both a membrane bound form and also a soluble aminopeptidase N.

Mouse anti Human CD13, clone WM15 inhibits infection of cells by human coronavirus ([Lachance *et al.* 1998](#)) but not hepatitis C virus ([Koutsoudakis *et al.* 2006](#)) and inhibits aminopeptidase N activity of the CD13 molecule ([Asmun *et al.* 1992](#)).

Flow Cytometry Use 10µl of the suggested working dilution to label 10⁶ cells or 100µl whole blood

References

1. Bradstock, K.F. *et al.* (1985) Myeloid progenitor surface antigen identified by monoclonal antibody. [Br J Haematol. 61 \(1\): 11-20.](#)
2. Favaloro, E.J. *et al.* (1988) Further characterization of human myeloid antigens (gp160,95; gp150; gp67): investigation of epitopic heterogeneity and non-haemopoietic

- distribution using panels of monoclonal antibodies belonging to CD-11b, CD-13 and CD-33. [Br J Haematol. 69 \(2\): 163-71.](#)
3. Favaloro, E.J. (1991) CD-13 (gp150; aminopeptidase-N): co-expression on endothelial and haemopoietic cells with conservation of functional activity. [Immunol Cell Biol. 69 \(Pt 4\): 253-60.](#)
 4. Favaloro, E.J. *et al.* (1993) The hepatobiliary disease marker serum alanine aminopeptidase predominantly comprises an isoform of the haematological myeloid differentiation antigen and leukaemia marker CD-13/gp150. [Clin Chim Acta. 220 \(1\): 81-90.](#)
 5. Favaloro, E.J. *et al.* (1993) CD13 (GP150; aminopeptidase-N): predominant functional activity in blood is localized to plasma and is not cell-surface associated. [Exp Hematol. 21 \(13\): 1695-701.](#)
 6. Gredmark, S. *et al.* (2004) Human Cytomegalovirus Induces Inhibition of Macrophage Differentiation by Binding to Human Aminopeptidase N/CD13 [J Immunol. 173: 4897-907](#)
 7. Thielitz, A. *et al.* (2004) Identification of extra- and intracellular alanyl aminopeptidases as new targets to modulate keratinocyte growth and differentiation. [Biochem Biophys Res Commun. 321 \(4\): 795-801.](#)
 8. Lassnig, C. *et al.* (2005) Development of a transgenic mouse model susceptible to human coronavirus 229E. [Proc Natl Acad Sci U S A. 102 \(23\): 8275-80.](#)
 9. Stolzing, A. *et al.* (2008) Age-related changes in human bone marrow-derived mesenchymal stem cells: consequences for cell therapies. [Mech Ageing Dev. 129: 163-73.](#)
 10. Negussie, A.H. *et al.* (2010) Synthesis and in vitro evaluation of cyclic NGR peptide targeted thermally sensitive liposome. [J Control Release. 143: 265-73.](#)
 11. Grzywacz, B. *et al.* (2011) Natural killer-cell differentiation by myeloid progenitors. [Blood. 117: 3548-58.](#)
 12. Tavoosidana, G. *et al.* (2011) Multiple recognition assay reveals prostasomes as promising plasma biomarkers for prostate cancer. [Proc Natl Acad Sci U S A. 108: 8809-14.](#)
 13. Silk, K.M. *et al.* (2012) Rapamycin conditioning of dendritic cells differentiated from human ES cells promotes a tolerogenic phenotype. [J Biomed Biotechnol. 2012:172420.](#)
 14. McCormack, E. *et al.* (2013) Multiplexed mAbs: a new strategy in preclinical time-domain imaging of acute myeloid leukemia. [Blood. 121 \(7\): e34-42.](#)
 15. Fiddler, C.A. *et al.* (2016) The Aminopeptidase CD13 Induces Homotypic Aggregation in Neutrophils and Impairs Collagen Invasion. [PLoS One. 11 \(7\): e0160108.](#)
 16. Chaturvedi, C.P. *et al.* (2018) Altered Expression of Hematopoiesis Regulatory Molecules in Lipopolysaccharide-Induced Bone Marrow Mesenchymal Stem Cells of Patients with Aplastic Anemia. [Stem Cells Int. 2018: 6901761.](#)
 17. Menon, R. *et al.* (2023) Human Induced Pluripotent Stem Cell-Derived Pericytes as Scalable and Editable Source to Study Direct Lineage Reprogramming Into Induced Neurons. [Cell Reprogram. 25 \(5\): 212-23.](#)
 18. Karpyuk, V. *et al.* (2019) Innovation-based Approach in Reconstruction of Reduced Jaw Alveolar Ridge Bone Using Cell Regeneration Technologies [Archiv Euromedica 9 \(2\) 147-55.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for

short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1270 10040
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M408988:221016'

Printed on 19 Jan 2024