

Datasheet: MCA1271A700

Description:	MOUSE ANTI HUMAN CD33:Alexa Fluor® 700		
Specificity:	CD33		
Format:	ALEXA FLUOR® 700		
Product Type:	Monoclonal Antibody		
Clone:	WM53		
Isotype:	lgG1		
Quantity:	100 TESTS/1ml		

# **Product Details**

# **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat - 1/5

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human				
Product Form	Purified IgG conjuga	0 - liquid			
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)		
	Alexa Fluor®700	702	723		
Preparation	Purified IgG prepare supernatant	ed by affinity chromatog	raphy on Protein G from tissue		
Buffer Solution	Phosphate buffered saline				
Preservative	0.09% sodium azide	e (NaN <sub>3</sub> )			
Stabilisers	1% bovine serum al				
Approx. Protein Concentrations	IgG concentration 0	.05 mg/ml			
Immunogen	Human AML cells				

UniProt:				
P20138 Related reagents				
Entrez Gene:				
945 CD33 Related reagents				
SIGLEC3				
AB_844496				
Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS myeloma cell line				

## **Specificity**

**Mouse anti Human CD33 antibody, clone WM53** recognizes the human CD33 cell surface glycoprotein. This antigen, considered to be specific for the myeloid lineage, has also been reported to be present on cells of lymphoid origin.

Mouse anti Human CD33 antibody, clone WM53 immunoprecipitates a protein of ~75 kDa from myeloid cells, a smaller protein of approximately 67 kDa has been observed in immunoprecipitates from lymphoid targets.

### Flow Cytometry

Use  $10\mu l$  of the suggested working dilution to label  $10^6$  cells or  $100\mu l$  whole blood.

### References

- 1. Favaloro, E.J. *et al.* (1988) Further characterization of human myeloid antigens (gp160,95; gp150; gp67): investigation of epitopic heterogeneity and non-haemopoietic distribution using panels of monoclonal antibodies belonging to CD-11b, CD-13 and CD-33. Br J Haematol. 69 (2): 163-71.
- 2. Favaloro, E.J. *et al.* (1993) Differential expression of surface antigens on activated endothelium. <u>Immunol Cell Biol. 71:571-81.</u>
- 3. Pietschmann, P. *et al.* (2000) Surface markers and transendothelial migration of dendritic cells from elderly subjects. Exp Gerontol. 35: 213-24.
- 4. Vamvakopoulos, J. *et al.* (2002) Genetic control of IL-1beta bioactivity through differential regulation of the IL-1 receptor antagonist. <u>Eur J Immunol. 32 (10): 2988-96.</u>
- 5. Vamvakopoulos, J.E. & Green, C. (2003) HMG-CoA reductase inhibition aborts functional differentiation and triggers apoptosis in cultured primary human monocytes: a potential mechanism of statin-mediated vasculoprotection. BMC Cardiovasc Disord. 3: 6.
- 6. Dahl C *et al.* (2004) Human mast cells express receptors for IL-3, IL-5 and GM-CSF; a partial map of receptors on human mast cells cultured *in vitro*. <u>Allergy. 59 (10): 1087-96.</u>
- 7. Lajaunias, F. *et al.* (2005) Constitutive repressor activity of CD33 on human monocytes requires sialic acid recognition and phosphoinositide 3-kinase-mediated intracellular signaling. <u>Eur J Immunol. 35: 243-51.</u>
- 8. Lin, C.W. *et al.* (2005) CD94 1A transcripts characterize lymphoblastic lymphoma/leukemia of immature natural killer cell origin with distinct clinical features. Blood. 106 (10): 3567-74.
- 9. Hernández-Caselles, T. *et al.* (2006) A study of CD33 (SIGLEC-3) antigen expression and function on activated human T and NK cells: two isoforms of CD33 are generated by alternative splicing. <u>J Leukoc Biol. 79: 46-58.</u>

- 10. Biedermann, B. *et al.* (2006) Analysis of the CD33-related siglec family reveals that Siglec-9 is an endocytic receptor expressed on subsets of acute myeloid leukemia cells and absent from normal hematopoietic progenitors. Leuk Res. 31: 211-20.
- 11. Yasukawa, T. *et al.* (2012) Simple detection of surface antigens on living cells by applying distinct cell positioning with negative dielectrophoresis. <u>Anal Chem. 84 (20):</u> 8830-6.
- 12. McCormack E *et al.* (2013) Multiplexed mAbs: a new strategy in preclinical time-domain imaging of acute myeloid leukemia. Blood. 121 (7): e34-42.
- 13. Hu, Z. *et al.* (2016) Self-assembled nanoparticles based on folic acid modified carboxymethyl chitosan conjugated with targeting antibody <u>J Wuhan Univ of Technol-Mater. Sci. Ed. 31 (2): 446-53.</u>
- 14. Hernández-Caselles T *et al.* (2019) CD33 (Siglec-3) Inhibitory Function: Role in the NKG2D/DAP10 Activating Pathway. <u>J Immunol Res. 2019: 6032141.</u>

### **Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

#### Guarantee

12 months from date of despatch

### Acknowledgements

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

# Health And Safety Information

Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1271A700">https://www.bio-rad-antibodies.com/SDS/MCA1271A700</a> 10041

### Regulatory

For research purposes only

# **Related Products**

## **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL: Alexa Fluor® 700 (MCA928A700)

## **Recommended Useful Reagents**

**HUMAN SEROBLOCK (BUF070A)** 

# **HUMAN SEROBLOCK (BUF070B)**

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M409017:221016'

## Printed on 23 Apr 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint