

Datasheet: MCA1271SBUV575

BATCH NUMBER 64765596

Description:	MOUSE ANTI HUMAN CD33:StarBright UltraViolet 575
Specificity:	CD33
Other names:	SIGLEC-3
Format:	StarBright UltraViolet 575
Product Type:	Monoclonal Antibody
Clone:	WM53
Isotype:	IgG1
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)Staining at lower temperatures generally requires longer incubation to reach antibody binding equilibrium, and this effect is more pronounced for StarBright Dyes due to their larger size. For optimum staining, Bio-Rad recommends incubating samples either overnight at 4°C or 1 hour at room temperature.

Target Species	Human		
Product Form	Purified IgG conjugated to StarBright UltraViolet 575 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	StarBright UltraViolet 575	340	569
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		

Preservative Stabilisers	0.09% sodium azide (NaN ₃) 1% bovine serum albumin 0.1% Pluronic F68 0.1% PEG 3350 0.05% Tween 20
Approx. Protein Concentrations	For information on the concentration of our StarBright Dye conjugated reagents please visit our FAQ page.
Immunogen	Human AML cells
External Database Links	UniProt: P20138 Related reagents Entrez Gene: 945 CD33 Related reagents
Synonyms	SIGLEC3
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1 myeloma cell line
Specificity	Mouse anti Human CD33 antibody, clone WM53, is a monoclonal antibody that recognizes CD33, also known as Siglec-3 or gp67. Clone WM53 detects CD33 in humans and has been validated in the following applications: flow cytometry, immunoprecipitation and western blotting. Clone WM53 immunoprecipitates a protein of ~75 kDa from myeloid cells, a smaller protein of approximately 67 kDa has been observed in immunoprecipitates from lymphoid targets. CD33 is a single pass type I membrane protein and member of the sialic acid-binding immunoglobulin-like lectin (Siglec) family. CD33 is expressed on myeloid progenitors, monocytes, granulocytes, dendritic cells and mast cells, but is absent on lymphocytes, normal platelets, hematopoietic stem cells and erythrocytes. Upon CD33 activation, the immunoreceptor tyrosine-based inhibition motif (ITIM) is phosphorylated by Src family kinases, resulting in recruitment of SH2 domain containing phosphatases and inhibition of cellular activation. As sialoglycans are present on almost all human cells, CD33 and other members of the siglec family have a role in discriminating self from non-self. Mouse anti human CD33 antibody, clone WM53, is also available as a recombinant antibody (TZA1271).
Flow Cytometry	Use 5µl of the suggested working dilution to label 0.5x10 ⁶ cells in 100µl. Best practices suggest a 5 min centrifugation at 6,000g prior to sample application.
References	1. Favaloro, E.J. <i>et al.</i> (1988) Further characterization of human myeloid antigens (gp160,95; gp150; gp67): investigation of epitopic heterogeneity and non-haemopoietic

- distribution using panels of monoclonal antibodies belonging to CD-11b, CD-13 and CD-33. [Br J Haematol. 69 \(2\): 163-71.](#)
2. Favaloro, E.J. *et al.* (1993) Differential expression of surface antigens on activated endothelium. [Immunol Cell Biol. 71:571-81.](#)
 3. Pietschmann, P. *et al.* (2000) Surface markers and transendothelial migration of dendritic cells from elderly subjects. [Exp Gerontol. 35: 213-24.](#)
 4. Vamvakopoulos, J. *et al.* (2002) Genetic control of IL-1beta bioactivity through differential regulation of the IL-1 receptor antagonist. [Eur J Immunol. 32 \(10\): 2988-96.](#)
 5. Vamvakopoulos, J.E. & Green, C. (2003) HMG-CoA reductase inhibition aborts functional differentiation and triggers apoptosis in cultured primary human monocytes: a potential mechanism of statin-mediated vasculoprotection. [BMC Cardiovasc Disord. 3: 6.](#)
 6. Dahl C *et al.* (2004) Human mast cells express receptors for IL-3, IL-5 and GM-CSF; a partial map of receptors on human mast cells cultured *in vitro*. [Allergy. 59 \(10\): 1087-96.](#)
 7. Lajaunias, F. *et al.* (2005) Constitutive repressor activity of CD33 on human monocytes requires sialic acid recognition and phosphoinositide 3-kinase-mediated intracellular signaling. [Eur J Immunol. 35: 243-51.](#)
 8. Lin, C.W. *et al.* (2005) CD94 1A transcripts characterize lymphoblastic lymphoma/leukemia of immature natural killer cell origin with distinct clinical features. [Blood. 106 \(10\): 3567-74.](#)
 9. Hernández-Caselles, T. *et al.* (2006) A study of CD33 (SIGLEC-3) antigen expression and function on activated human T and NK cells: two isoforms of CD33 are generated by alternative splicing. [J Leukoc Biol. 79: 46-58.](#)
 10. Biedermann, B. *et al.* (2006) Analysis of the CD33-related siglec family reveals that Siglec-9 is an endocytic receptor expressed on subsets of acute myeloid leukemia cells and absent from normal hematopoietic progenitors. [Leuk Res. 31: 211-20.](#)
 11. Yasukawa, T. *et al.* (2012) Simple detection of surface antigens on living cells by applying distinct cell positioning with negative dielectrophoresis. [Anal Chem. 84 \(20\): 8830-6.](#)
 12. McCormack E *et al.* (2013) Multiplexed mAbs: a new strategy in preclinical time-domain imaging of acute myeloid leukemia. [Blood. 121 \(7\): e34-42.](#)
 13. Hu, Z. *et al.* (2016) Self-assembled nanoparticles based on folic acid modified carboxymethyl chitosan conjugated with targeting antibody [J Wuhan Univ of Technol-Mater. Sci. Ed. 31 \(2\): 446-53.](#)
 14. Hernández-Caselles T *et al.* (2019) CD33 (Siglec-3) Inhibitory Function: Role in the NKG2D/DAP10 Activating Pathway. [J Immunol Res. 2019: 6032141.](#)

Storage

This product is shipped at ambient temperature.
Store at +4°C.
DO NOT FREEZE.
This product should be stored undiluted.

Guarantee

12 months from date of despatch

Acknowledgements

This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts

Health And Safety Information

Material Safety Datasheet documentation #20471 available at:
<https://www.bio-rad-antibodies.com/SDS/MCA1271SBUV575>

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M451563:260710'

Printed on 21 Aug 2026