# Datasheet: MCA1271SBUV665 BATCH NUMBER 100006134

Description:	MOUSE ANTI HUMAN CD33:StarBright UltraViolet 665			
Specificity:	CD33			
Format:	StarBright UltraViolet 665			
Product Type:	Monoclonal Antibody			
Clone:	WM53			
Isotype:	lgG1			
Quantity:	100 TESTS/0.5ml			

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes N	0	Not Determined	Suggested Dilution	
	Flow Cytometry	•			Neat	
	Where this product ha necessarily exclude its a guide only. It is recon system using appropria	s use in such pro mmended that th	cedure e user	es. Suggested workin titrates the product	ng dilutions are given as	
Target Species	Human					
Product Form	Purified IgG conjugate	d to StarBright U	ItraVio	let 665 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (	nm) l	Emission Max (nm)		
	StarBright UltraViolet 665	340		684		
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant					
Buffer Solution	Phosphate buffered saline					
Preservative Stabilisers	0.09% Sodium Azide ( 1% Bovine Serum Albr 0.1% Pluronic F68 0.1% PEG 3350	•••				

	0.05% Tween 20
Immunogen	Human AML cells
External Database Links	UniProt: <u>P20138</u> <u>Related reagents</u> Entrez Gene: <u>945</u> CD33 <u>Related reagents</u>
Synonyms	SIGLEC3
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1 myeloma cell line
Specificity	<b>Mouse anti Human CD33 antibody, clone WM53</b> recognizes the human CD33 cell surface glycoprotein. This antigen, considered to be specific for the myeloid lineage, has also been reported to be present on cells of lymphoid origin.
	Mouse anti Human CD33 antibody, clone WM53 immunoprecipitates a protein of ~75 kDa from myeloid cells, a smaller protein of approximately 67 kDa has been observed in immunoprecipitates from lymphoid targets.
Flow Cytometry	Use 5ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
References	<ol> <li>Favaloro, E.J. <i>et al.</i> (1987) Characterization of monoclonal antibodies to the human myeloid-differentiation antigen, gp67 (CD-33). <u>Dis Markers. 5 (4): 215-25.</u></li> <li>Favaloro, E.J. <i>et al.</i> (1988) Further characterization of human myeloid antigens (gp160,95; gp150; gp67): investigation of epitopic heterogeneity and non-haemopoietic distribution using panels of monoclonal antibodies belonging to CD-11b, CD-13 and CD-33. <u>Br J Haematol. 69 (2): 163-71.</u></li> <li>Hernández-Caselles, T. <i>et al.</i> (2006) A study of CD33 (SIGLEC-3) antigen expression and function on activated human T and NK cells: two isoforms of CD33 are generated by alternative splicing. <u>J Leukoc Biol. 79: 46-58.</u></li> <li>Biedermann, B. <i>et al.</i> (2006) Analysis of the CD33-related siglec family reveals that Siglec-9 is an endocytic receptor expressed on subsets of acute myeloid leukemia cells and absent from normal hematopoietic progenitors. <u>Leuk Res. 31: 211-20.</u></li> <li>Lajaunias, F. <i>et al.</i> (2005) Constitutive repressor activity of CD33 on human monocytes requires sialic acid recognition and phosphoinositide 3-kinase-mediated intracellular signaling. <u>Eur J Immunol. 35: 243-51.</u></li> <li>Pietschmann, P. <i>et al.</i> (2000) Surface markers and transendothelial migration of dendritic cells from elderly subjects. <u>Exp Gerontol. 35: 213-24.</u></li> <li>Favaloro, E.J. <i>et al.</i> (2012) Simple detection of surface antigens on activated endothelium. <u>Immunol Cell Biol. 71:571-81.</u></li> <li>Yasukawa, T. <i>et al.</i> (2012) Simple detection of surface antigens on living cells by applying distinct cell positioning with negative dielectrophoresis. <u>Anal Chem. 84 (20):</u></li> </ol>

	<u>8830-6.</u>
	9. Hu, Z. et al. (2016) Self-assembled nanoparticles based on folic acid modified
	carboxymethyl chitosan conjugated with targeting antibody <u>J Wuhan Univ of Technol-</u>
	<u>Mater. Sci. Ed. 31 (2): 446-53.</u>
	10. Dahl C et al. (2004) Human mast cells express receptors for IL-3, IL-5 and GM-CSF; a
	partial map of receptors on human mast cells cultured <i>in vitro</i> . <u>Allergy. 59 (10): 1087-96.</u>
	11. Vamvakopoulos, J.E. & Green, C. (2003) HMG-CoA reductase inhibition aborts
	functional differentiation and triggers apoptosis in cultured primary human monocytes: a
	potential mechanism of statin-mediated vasculoprotection. <u>BMC Cardiovasc Disord. 3: 6.</u>
	12. Vamvakopoulos, J. et al. (2002) Genetic control of IL-1beta bioactivity through
	differential regulation of the IL-1 receptor antagonist. Eur J Immunol. 32 (10): 2988-96.
	13. Lin, C.W. et al. (2005) CD94 1A transcripts characterize lymphoblastic
	lymphoma/leukemia of immature natural killer cell origin with distinct clinical features.
	Blood. 106 (10): 3567-74.
	14. McCormack E <i>et al.</i> (2013) Multiplexed mAbs: a new strategy in preclinical
	time-domain imaging of acute myeloid leukemia. <u>Blood. 121 (7): e34-42.</u>
	15. Hernández-Caselles T et al. (2019) CD33 (Siglec-3) Inhibitory Function: Role in the
	NKG2D/DAP10 Activating Pathway. <u>J Immunol Res. 2019: 6032141.</u>
Storage	Store at +4°C. DO NOT FREEZE.
	This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety	Material Safety Datasheet documentation #20471 available at:
Information	https://www.bio-rad-antibodies.com/SDS/MCA1271SBUV665
	20471
Regulatory	For research purposes only

## **Related Products**

### **Recommended Useful Reagents**

### HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	id.com	Email: antibody_sales_uk@bio-ra	id.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M393385:211221'

#### Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint