

Datasheet: MCA1271SBY575

Description:	MOUSE ANTI HUMAN CD33:StarBright Yellow 575
Specificity:	CD33
Format:	StarBright Yellow 575
Product Type:	Monoclonal Antibody
Clone:	WM53
Isotype:	IgG1
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to StarBright Yellow 575 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	StarBright Yellow 575	548	579
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		
	0.1% Pluronic F68		
	0.1% PEG 3350		
	0.05% Tween 20		

Immunogen	Human AML cells
External Database Links	<p>UniProt: P20138 Related reagents</p> <p>Entrez Gene: 945 CD33 Related reagents</p>
Synonyms	SIGLEC3
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1 myeloma cell line
Specificity	<p>Mouse anti Human CD33 antibody, clone WM53 recognizes the human CD33 cell surface glycoprotein. This antigen, considered to be specific for the myeloid lineage, has also been reported to be present on cells of lymphoid origin.</p> <p>Mouse anti Human CD33 antibody, clone WM53 immunoprecipitates a protein of ~75 kDa from myeloid cells, a smaller protein of approximately 67 kDa has been observed in immunoprecipitates from lymphoid targets.</p>
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 ⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
References	<ol style="list-style-type: none"> 1. Favaloro, E.J. <i>et al.</i> (1988) Further characterization of human myeloid antigens (gp160,95; gp150; gp67): investigation of epitopic heterogeneity and non-haemopoietic distribution using panels of monoclonal antibodies belonging to CD-11b, CD-13 and CD-33. Br J Haematol. 69 (2): 163-71. 2. Favaloro, E.J. <i>et al.</i> (1993) Differential expression of surface antigens on activated endothelium. Immunol Cell Biol. 71:571-81. 3. Pietschmann, P. <i>et al.</i> (2000) Surface markers and transendothelial migration of dendritic cells from elderly subjects. Exp Gerontol. 35: 213-24. 4. Vamvakopoulos, J. <i>et al.</i> (2002) Genetic control of IL-1beta bioactivity through differential regulation of the IL-1 receptor antagonist. Eur J Immunol. 32 (10): 2988-96. 5. Vamvakopoulos, J.E. & Green, C. (2003) HMG-CoA reductase inhibition aborts functional differentiation and triggers apoptosis in cultured primary human monocytes: a potential mechanism of statin-mediated vasculoprotection. BMC Cardiovasc Disord. 3: 6. 6. Dahl C <i>et al.</i> (2004) Human mast cells express receptors for IL-3, IL-5 and GM-CSF; a partial map of receptors on human mast cells cultured <i>in vitro</i>. Allergy. 59 (10): 1087-96. 7. Lajaunias, F. <i>et al.</i> (2005) Constitutive repressor activity of CD33 on human monocytes requires sialic acid recognition and phosphoinositide 3-kinase-mediated intracellular signaling. Eur J Immunol. 35: 243-51. 8. Lin, C.W. <i>et al.</i> (2005) CD94 1A transcripts characterize lymphoblastic lymphoma/leukemia of immature natural killer cell origin with distinct clinical features. Blood. 106 (10): 3567-74. 9. Hernández-Caselles, T. <i>et al.</i> (2006) A study of CD33 (SIGLEC-3) antigen expression and function on activated human T and NK cells: two isoforms of CD33 are generated by

alternative splicing. [J Leukoc Biol. 79: 46-58.](#)

10. Biedermann, B. *et al.* (2006) Analysis of the CD33-related siglec family reveals that Siglec-9 is an endocytic receptor expressed on subsets of acute myeloid leukemia cells and absent from normal hematopoietic progenitors. [Leuk Res. 31: 211-20.](#)

11. Yasukawa, T. *et al.* (2012) Simple detection of surface antigens on living cells by applying distinct cell positioning with negative dielectrophoresis. [Anal Chem. 84 \(20\): 8830-6.](#)

12. McCormack E *et al.* (2013) Multiplexed mAbs: a new strategy in preclinical time-domain imaging of acute myeloid leukemia. [Blood. 121 \(7\): e34-42.](#)

13. Hu, Z. *et al.* (2016) Self-assembled nanoparticles based on folic acid modified carboxymethyl chitosan conjugated with targeting antibody [J Wuhan Univ of Technol-Mater. Sci. Ed. 31 \(2\): 446-53.](#)

14. Hernández-Caselles T *et al.* (2019) CD33 (Siglec-3) Inhibitory Function: Role in the NKG2D/DAP10 Activating Pathway. [J Immunol Res. 2019: 6032141.](#)

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
----------------	--

Guarantee	12 months from date of despatch
------------------	---------------------------------

Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
-------------------------	---

Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA1271SBY575 20471
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M413704:221125'

Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)