

Datasheet: MCA1298F

Description:	MOUSE ANTI HUMAN CD79a:FITC
Specificity:	CD79a
Other names:	MB-1
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	ZL7-4
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Antibody purified from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% sodium azide (NaN ₃)		
Stabilisers	1% bovine serum albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		

Immunogen IgM complex isolated from Daudi cells.

External Database

Links

UniProt:

[P11912](#) [Related reagents](#)

Entrez Gene:

[973](#) CD79A [Related reagents](#)

Synonyms

IGA, MB1

RRID

AB_321768

Fusion Partners

Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.

Specificity

Mouse anti Human CD79a antibody, clone ZL7-4 recognizes the human B-cell antigen receptor complex-associated protein alpha chain, also known as MB-1 membrane glycoprotein or CD79a. clone ZL7-4 reacts with CD79a positive cells by flow cytometry and with CD79a in an ELISA specific for a fusion protein of CD79a-Fc.

Mouse anti Human CD79a antibody, clone ZL7-4 has been reported to be useful in distinguishing B-CLL from mantle cell lymphoma in flow cytometric assays ([Bell et al. 1999](#)).

Mouse anti Human CD79a antibody, clone ZL7-4 has been reported to be suitable for Immunohistochemistry on frozen and pre-treated paraffin sections, but does exhibit epithelial staining.

Mouse anti Human CD79a antibody, clone ZL7-4 has been reported to induce phosphorylation of syk kinase ([Lanham et al. 2003](#)).

Flow Cytometry

Use 10µl of the suggested working dilution to label 10⁶ cells or 100µl whole blood We recommend incubation times of at least 30 minutes with this antibody.

References

1. Zhang, L. *et al.* (1995) The development of anti-CD79 monoclonal antibodies for treatment of B-cell neoplastic disease. [Therapeutic Immunology 2:191-202](#)
2. Bell, P.B. *et al.* (1999) CD79a detected by ZL7.4 separates chronic lymphocytic leukemia from mantle cell lymphoma in the leukemic phase. [Cytometry. 38 \(3\): 102-5.](#)
3. Cragg, M.S. *et al.* (2002) The alternative transcript of CD79b is overexpressed in B-CLL and inhibits signaling for apoptosis. [Blood. 100: 3068-76.](#)
4. Lanham, S. *et al.* (2003) Differential signaling via surface IgM is associated with VH gene mutational status and CD38 expression in chronic lymphocytic leukemia. [Blood. 101 \(3\): 1087-93.](#)
5. Allsup, D.J. *et al.* (2005) B-cell receptor translocation to lipid rafts and associated signaling differ between prognostically important subgroups of chronic lymphocytic leukemia. [Cancer Res. 65: 7328-37.](#)
6. Rahemtullah, A. *et al.* (2008) CD20+ T-cell lymphoma: clinicopathologic analysis of 9

cases and a review of the literature. [Am J Surg Pathol. 32 \(11\): 1593-607.](#)

7. Luger, D. *et al.* (2013) Expression of the B-cell receptor component CD79a on immature myeloid cells contributes to their tumor promoting effects. [PLoS One. 8 \(10\): e76115.](#)

8. Vendel, A.C. *et al* (2009) B and T lymphocyte attenuator regulates B cell receptor signaling by targeting Syk and BLNK [J Immunol. 182: 1509-17.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1298F>
10041

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M410381:221028'

Printed on 24 Apr 2025

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)