

Datasheet: MCA1302G

### **BATCH NUMBER 154985**

Description:	RAT ANTI MOUSE INTERLEUKIN-10		
Specificity:	IL-10		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	JES5-2A5		
Isotype:	lgG1		
Quantity:	0.5 mg		

# **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	•			1/10 - 1/100
Immunohistology - Frozen	•			
Immunohistology - Paraffin				
ELISA				1/200 - 1/2000
Immunoprecipitation				
Western Blotting	•			1/1000

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) Membrane permeabilization is required for this application. Bio-Rad recommends the use of Leucoperm (BUF09) for this purpose.

Target Species	Mouse
Species Cross Reactivity	Reacts with: Goat  N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture

supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Recombinant Mouse IL-10.
External Database Links	UniProt: P18893 Related reagents
	Entrez Gene:  16153 II10 Related reagents
Synonyms	II-10
RRID	AB_2125093
Fusion Partners	Spleen cells from immunized rats were fused with cells of the mouse P3.X63 Ag8.653 myeloma cell line.
Specificity	Rat anti Mouse Interleukin-10 antibody, clone JES5-2A5 recognizes mouse Cytokine synthesis inhibitory factor (CSIF), also known as interleukin-10 (IL-10). IL-10 is produced by Ly-1(CD5) expressing B cells (O'Garra et al. 1992) and a subset of stimulated T cells (Asseman et al. 1999).
	IL-10 has a wide range of biological properties, including the inhibition of macrophage and T helper cell mediated cytokine synthesis, and the suppression of the delayed hypersensitivity response.
	Rat anti Mouse Interleukin-10 antibody, clone JES5-2A5 is reported to neutralize the activity of murine IL-10 (Weber-Nordt et al. 1994).
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.
References	<ol> <li>Abrams, J.S. <i>et al.</i> (1992) Strategies of anti-cytokine monoclonal antibody development: immunoassay of IL-10 and IL-5 in clinical samples. <u>Immunological Review 127: 5-24.</u></li> <li>Sander, B. <i>et al.</i> (1993) Similar frequencies and kinetics of cytokine producing cells in murine peripheral blood and spleen. Cytokine detection by immunoassay and intracellular immunostaining. <u>J Immunol Methods. 166 (2): 201-14.</u></li> <li>Nishigori, C. <i>et al.</i> (1998) HindIII liposomes suppress delayed-type hypersensitivity responses <i>in vivo</i> and induce epidermal IL-10 <i>in vitro</i>. <u>J Immunol. 161 (6): 2684-91.</u></li> </ol>

- 4. Weber-nordt, R.M. *et al.* (1994) Lipopolysaccharide-dependent induction of IL-10 receptor expression on murine fibroblasts. J Immunol. 153 (8): 3734-44.
- 5. Rennick, D. *et al.* (1995) Cofactors are essential for stem cell factor-dependent growth and maturation of mast cell progenitors: comparative effects of interleukin-3 (IL-3), IL-4, IL-10, and fibroblasts. <u>Blood. 85: 57-65.</u>
- 6. Biswas, S.K. *et al.* (2006) A distinct and unique transcriptional program expressed by tumor-associated macrophages (defective NF-kappaB and enhanced IRF-3/STAT1 activation). Blood. 107: 2112-22.
- 7. Gaupp, S. *et al.* (2008) Amelioration of experimental autoimmune encephalomyelitis in IL-4Ralpha-/- mice implicates compensatory up-regulation of Th2-type cytokines. <u>Am J Pathol.</u> 173: 119-29.
- 8. Gonzalez-Juarrero, M. *et al.* (2013) Polar Lipids of *Burkholderia pseudomallei* Induce Different Host Immune Responses. PLoS One. 8: e80368.
- 9. Higgins, J.L. *et al.* (2018) Cell mediated immune response in goats after experimental challenge with the virulent *Brucella melitensis* strain 16M and the reduced virulence strain Rev. 1. Vet Immunol Immunopathol. 202: 74-84.
- 10. Hammond, J.W. *et al.* (2018) HIV Tat causes synapse loss in a mouse model of HIV-associated neurocognitive disorder that is independent of the classical complement cascade component C1q. <u>Glia. 66 (12): 2563-74.</u>

### **Storage**

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1302G">https://www.bio-rad-antibodies.com/SDS/MCA1302G</a> 10040
Regulatory	For research purposes only

## Related Products

# **Recommended Secondary Antibodies**

Rabbit Anti Rat IgG (STAR16...)

DyLight®800

Rabbit Anti Rat IgG (STAR17...)

Goat Anti Rat IgG (STAR72...)

Goat Anti Rat IgG (STAR69...)

Goat Anti Rat IgG (STAR73...)

RPE

Rabbit Anti Rat IgG (STAR21...)

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) DyLight®550, DyLight®650, DyLight®800

Goat Anti Rat IgG (STAR131...) Alk. Phos., Biotin

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739 Fax: +49 (0) 89 8090 95 50 To

Email: antibody\_sales\_us@bio-rad.com Email: antibody\_sales\_uk@bio-rad.com Email: antibody\_sales\_de@bio-rad.com

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M365114:200529'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint