

Datasheet: MCA1304G

BATCH NUMBER 164020

Description: RAT ANTI MOUSE INTERLEUKIN		
Specificity:	IL-4	
Format:	Purified	
Product Type:	Monoclonal Antibody	
Clone:	BVD4-1D11	
Isotype:	lgG2b	
Quantity:	0.5 mg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	•			1/10 - 1/50
Immunohistology - Frozen	•			
Immunohistology - Paraffin				
ELISA	•			1ug/ml - 5ug/ml
Immunoprecipitation				
Western Blotting	•			1ug/ml - 5ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) Membrane permeabilization is required for this application. The use of Leucoperm (Product Code <u>BUF09</u>) is recommended for this purpose.

Target Species	Mouse	
Species Cross Reactivity	Does not react with:Rat	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on protein G supernatant	from tissue culture
Buffer Solution	Phosphate buffered saline	

Preservative Stabilisers	0.09% sodium azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Recombinant Mouse IL-4.
External Database Links	UniProt: P07750 Related reagents Entrez Gene: 16189 II4 Related reagents
Synonyms	11-4
RRID	AB_322398
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse P3.X63 Ag8.653 myeloma cell line.
Specificity	Rat anti Mouse Interleukin-4 antibody, clone BVD4-1D11 recognizes natural and recombinant mouse IL-4.
	Clone BVD4-1D11 is reported to neutralize IL-4 activity (Abrams et al., 1992).
Flow Cytometry	Use 10µl of the suggested working dilution to label 1x10 ⁶ cells in 100µl
References	 Abrams, J.S. <i>et al.</i> (1992) Strategies of anti-cytokine monoclonal antibody development immunoassay of IL-10 and IL-5 in clinical samples. Immunol Rev. 127: 5-24. Sander, B. <i>et al.</i> (1993) Similar frequencies and kinetics of cytokine producing cells in murine peripheral blood and spleen. Cytokine detection by immunoassay and intracellular immunostaining. J Immunol Methods. 166 (2): 201-14. Hochrein, H. <i>et al.</i> (2000) Interleukin (IL)-4 is a major regulatory cytokine governing bioactive IL-12 production by mouse and human dendritic cells. J Exp Med. 192 (6): 823-33. Finkelman, F.D. <i>et al.</i> (2000) Stat6 regulation of <i>in vivo</i> IL-4 responses. J Immunol. 16: 2303-10. Hochrein, H. <i>et al.</i> (2000) Interleukin (IL)-4 is a major regulatory cytokine governing bioactive IL-12 production by mouse and human dendritic cells. J Exp Med. 192: 823-33. Urban, J.F. Jr. <i>et al.</i> (2000) Stat6 signaling promotes protective immunity against Trichinella spiralis through a mast cell- and T cell-dependent mechanism. J Immunol. 16: 2046-52. Yang, Y.F. <i>et al.</i> (2000) Requirement for IFN-gamma in IL-12 production induced by collaboration between v(alpha)14(+) NKT cells and antigen-presenting cells. Int Immunol.

12: 1669-75.

8. Pechová, J. *et al.* (2004) Effect of tick salivary gland extract on the cytokine production by mouse epidermal cells. Folia Parasitol (Praha). 51: 367-72.

9. Roy, E. *et al.* (2008) Beneficial effect of anti-interleukin-4 antibody when administered in a murine model of tuberculosis infection. <u>Tuberculosis</u> (Edinb). 88 (3): 197-202.

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1304G 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR69...) FITC

Goat Anti Rat IgG (STAR73...) RPE

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) <u>DyLight®550</u>, <u>DyLight®650</u>, <u>DyLight®800</u>

Rabbit Anti Rat IgG (STAR21...) HRP

Rabbit Anti Rat IgG (STAR16...) <u>DyLight®800</u>
Goat Anti Rat IgG (STAR131...) <u>Alk. Phos., Biotin</u>

Rabbit Anti Rat IgG (STAR17...) <u>FITC</u>
Goat Anti Rat IgG (STAR72...) <u>HRP</u>

North & South Tel: +1 800 265 7376 Wor America Fax: +1 919 878 3751

Worldwide Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M409123:221017'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint