

Datasheet: MCA1322A647T BATCH NUMBER 1709

Description:	RAT ANTI MOUSE CD204:Alexa Fluor® 647		
Specificity:	CD204		
Other names:	SCAVENGER RECEPTOR TYPE I/II		
Format:	ALEXA FLUOR® 647		
Product Type:	Monoclonal Antibody		
Clone:	2F8		
lsotype:	lgG2b		
Quantity:	25 TESTS/0.25ml		

Product Details

Applications	This product has been derived from testing wi communications from t information. For genera rad-antibodies.com/pro	tions or personal ndicated for further			
		Yes No	Not Determined	Suggested Dilution	
	Flow Cytometry			Neat - 1/10	
	Where this antibody ha	as not been tested	or use in a particular teo	chnique this does not	
	•	nmended that the u	lures. Suggested workin iser titrates the antibody e controls.	•	
Target Species	Mouse				
Species Cross	Reacts with: Pig, Chan	nel catfish			
Reactivity	reactivity is derived from	m testing within ou	itions may vary betweer laboratories, peer-revie tors. Please refer to refe	wed publications or	
Product Form	Purified IgG conjugated to Alexa Fluor® 647 - liquid				
Max Ex/Em	Fluorophore	Excitation Max (nm) Emission Max (nm)		
	Alexa Fluor®647	650	665		
Preparation	Purified IgG prepared t supernatant	by affinity chromato	graphy on Protein G froi	m tissue culture	

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml
Immunogen	Raw 264 cell line
External Database Links	UniProt: P30204 Related reagents Entrez Gene: 20288 Msr1 Related reagents
Synonyms	Scvr
RRID	AB_2148251
Fusion Partners	Spleen cells from immunised AO rats were fused with cells of the Y3 rat myeloma cell line
Specificity	Rat anti Mouse CD204 antibody, clone 2F8 recognizes the murine scavenger receptor class A (SR-A), type I and II, also known as CD204. CD204 is expressed by tissue macrophages and functions both as an endocytic receptor for lipoproteins and as an adhesion receptor for macrophages binding to ligand rich tissues e.g. atherosclerotic lesions. Rat anti Mouse CD204 antibody, clone 2F8 inhibits the uptake of acetylated low-density lipoproteins and also inhibits divalent cation independent adhesion (Fraser <i>et al.</i> 1993).
	Rat anti Mouse CD204 antibody, clone 2F8 recognizes an epitope within SRA that is polymorphic in the SRA from C57BL/6 mice. Rat anti Mouse CD204 antibody, clone 2F8 is therefore unsuitable for use with the C57BL/6 mouse strain (<u>Daugherty <i>et al.</i> 2000</u>).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
	The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR (<u>BUF041A/B</u>).
References	 Fraser, I. <i>et al.</i> (1993) Divalent cation-independent macrophage adhesion inhibited by monoclonal antibody to murine scavenger receptor. <u>Nature. 364 (6435): 343-6.</u> de Villiers, W.J. <i>et al.</i> (1994) Macrophage-colony-stimulating factor selectively enhances macrophage scavenger receptor expression and function. <u>J Exp Med. 180 (2): 705-9.</u> Hughes, D.A. <i>et al.</i> (1995) Murine macrophage scavenger receptor: in vivo expression and function as receptor for macrophage adhesion in lymphoid and non-lymphoid organs. <u>Eur J Immunol. 25 (2): 466-73.</u> Gordon, S. (1995) The macrophage. <u>Bioessays. 17 (11): 977-86.</u>

5. Hughes, D.A. *et al.* (1994) Murine M phi scavenger receptor: adhesion function and expression. <u>Immunol Lett. 43 (1-2): 7-14.</u>

6. Aid, S. *et al.* (2008) Neuroinflammatory response to lipopolysaccharide is exacerbated in mice genetically deficient in cyclooxygenase-2. <u>J Neuroinflammation. 5: 17.</u>

7. Daugherty, A. *et al.* (2000) Polymorphism of class A scavenger receptors in C57BL/6 mice. J Lipid Res. 41 (10): 1568-77.

8. Moldenhauer, L.M. *et al.* (2010) GM-CSF is an essential regulator of T cell activation competence in uterine dendritic cells during early pregnancy in mice. <u>J Immunol. 185 (11):</u> 7085-96.

9. Luechtenborg, B. *et al.* (2008) Function of scavenger receptor class A type I/II is not important for smooth muscle foam cell formation. <u>Eur J Cell Biol. 87: 91-9.</u>

10. Sever-Chroneos, Z. *et al.* (2011) Surfactant Protein A (SP-A)-mediated Clearance of *Staphylococcus aureus* Involves Binding of SP-A to the Staphylococcal Adhesin Eap and the Macrophage Receptors SP-A Receptor 210 and Scavenger Receptor Class A. J Biol Chem. 286: 4854-70.

11. Yang, C.N. *et al.* (2011) Mechanism mediating oligomeric Aβ clearance by naïve primary microglia. <u>Neurobiol Dis. 42 (3): 221-30.</u>

12. Hald, A. *et al.* (2011) MMP9 is protective against lethal inflammatory mass lesions in the mouse colon. <u>Dis Model Mech. 4: 212-27.</u>

Swain, S.D. *et al.* (2011) *Pneumocystis* infection in an immunocompetent host can promote collateral sensitization to respiratory antigens. <u>Infect Immun. 79 (5): 1905-14.</u>
 Nikolic, D. *et al.* (2011) SR-A ligand and M-CSF dynamically regulate SR-A expression

and function in primary macrophages via p38 MAPK activation. <u>BMC Immunol. 12: 37.</u>

15. Zaynagetdinov, R *et al.* (2011) A critical role for macrophages in promotion of urethane-induced lung carcinogenesis. J Immunol. 187 (11): 5703-11.

16. Kaur, H. *et al.* (2003) Identification of a scavenger receptor homologue on nonspecific cytotoxic cells and evidence for binding to oligodeoxyguanosine. <u>Fish Shellfish Immunol.</u> <u>15: 169-81.</u>

17. Kaur, H. *et al.* (2004) Single-base oligodeoxyguanosine-binding proteins on nonspecific cytotoxic cells: identification of a new class of pattern-recognition receptors. <u>Scand J Immunol. 60: 238-48.</u>

18. Koronyo Y *et al.* (2015) Therapeutic effects of glatiramer acetate and grafted CD115+ monocytes in a mouse model of Alzheimer's disease. <u>Brain. 138 (Pt 8): 2399-422.</u>

19. Nielsen, B.S. *et al.* (2008) Matrix metalloproteinase 13 is induced in fibroblasts in polyomavirus middle T antigen-driven mammary carcinoma without influencing tumor progression. <u>PLoS One. 3 (8): e2959.</u>

20. Tao, J. *et al.* (2015) CIC-3 deficiency prevents atherosclerotic lesion development in ApoE^{-/-} mice. J Mol Cell Cardiol. 87: 237-247.

21. Prins, J.R. *et al.* (2015) Unstable Foxp3+ regulatory T cells and altered dendritic cells are associated with lipopolysaccharide-induced fetal loss in pregnant interleukin 10-deficient mice. <u>Biol Reprod. 93 (4): 95.</u>

22. Almholt, K. *et al.* (2015) Spontaneous lung and lymph node metastasis in transgenic breast cancer is independent of the urokinase receptor uPAR. <u>Clin Exp Metastasis. 32 (6):</u> 543-54.

23. Verheijden S *et al.* (2015) Identification of a chronic non-neurodegenerative microglia activation state in a mouse model of peroxisomal β -oxidation deficiency. <u>Glia. 63 (9):</u> <u>1606-20.</u>

	 24. Kokubu, Y. <i>et al.</i> (2016) Induction of protumoral CD11c(high) macrophages by glioma cancer stem cells through GM-CSF. <u>Genes Cells. 21 (3): 241-51.</u> 25. Sapkota, M. <i>et al.</i> (2016) Malondialdehyde-Acetaldehyde-Adducted Surfactant Protein Alters Macrophage Functions Through Scavenger Receptor A. <u>Alcohol Clin Exp Res. 40</u> (12): 2563-2572. 26. Fujiwara, Y. <i>et al.</i> (2016) Onionin A, a sulfur-containing compound isolated from onions, impairs tumor development and lung metastasis by inhibiting the protumoral and immunosuppressive functions of myeloid cells. <u>Mol Nutr Food Res. Jul 9. [Epub ahead of print]</u> 27. Tsay, H.J. <i>et al.</i> (2016) Identifying N-linked glycan moiety and motifs in the cysteine-rich domain critical for N-glycosylation and intracellular trafficking of SR-AI and MARCO. <u>J Biomed Sci. 23: 27.</u> 28. Horlad, H. <i>et al.</i> (2013) Corosolic acid impairs tumor development and lung metastasis by inhibiting the immunosuppressive activity of myeloid-derived suppressor cells. <u>Mol Nutr Food Res. 57 (6): 1046-54.</u> 29. Shiau, D.J. <i>et al.</i> (2020) Hepatocellular carcinoma-derived high mobility group box 1 triggers M2 macrophage polarization via a TLR2/NOX2/autophagy axis. <u>Sci Rep. 10 (1): 13582.</u> 30. Tian, L.X. <i>et al.</i> (2021) Cytochrome P450 1A1 enhances inflammatory responses and impedes phagocytosis of bacteria in macrophages during sepsis. <u>Cell Commun Signal. 18 (1): 70.</u> 31. Wang, Y. <i>et al.</i> (2021) Scavenger receptor A1 participates in uptake of <i>Leptospira interrogans</i> serovar Autumnalis strain 56606v and inflammation in mouse macrophages.
Storage	Emerg Microbes Infect. : 1-39. Store at +4°C or at -20°C if preferred.
	This product should be stored undiluted.
	Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.
	Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Acknowledgements	This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

			neet documentation #10 tibodies.com/SDS/MCA		e at:
y	For researc	ch purposes	s only		
		Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
Email: antibody_sຄ	ales_us@bio-rad	l.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com
F	/ Fel: +1 800 265 73 Fax: +1 919 878 3	10041 For researc	10041 For research purposes Fel: +1 800 265 7376 Worldwide	Integr.//www.blog-rad-antibodies.com/SDS/MCA 10041 For research purposes only Fel: +1 800 265 7376 Worldwide Tel: +44 (0)1865 852 700 Fax: +1 919 878 3751	Intps://www.bio-rad-antibodies.com/bb/s/MCA1522A0471 10041 For research purposes only Fel: +1 800 265 7376 Worldwide Tel: +44 (0)1865 852 700 Europe Fax: +1 919 878 3751

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint