

Datasheet: MCA1322EL

Description:	RAT ANTI MOUSE CD204:Low Endotoxin		
Specificity:	CD204		
Other names:	SCAVENGER RECEPTOR TYPE I/II		
Format:	Low Endotoxin		
Product Type:	Monoclonal Antibody		
Clone:	2F8		
Isotype:	lgG2b		
Quantity:	0.5 mg		

Product Details

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry	-			1/50 - 1/100	
	Immunohistology - Frozen	-				
	(1)	_				
	Immunohistology - Paraffin		•			
	Immunohistology - Resin		-			
	ELISA	-				
	Immunoprecipitation	-				
	Western Blotting (2)	-				
	Functional Assays	-				
	Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls. (1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of					
acetone fixation for frozen sections.						
	(2)This product recognises CD204 in J774 cells under non-reduced conditi				duced conditions only.	
Target Species	Mouse					
Species Cross	Reacts with: Pig, Channe	el catfish				
Reactivity	N.B. Antibody reactivity and working conditions may vary between species. Cross					
	reactivity is derived from		-		•	
	reastivity is derived norm	coung w				

	personal communications from the originators. Please refer to further information.	references indicated for
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein G supernatant	from tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	None present	
Carrier Free	Yes	
Endotoxin Level	<0.01 EU/µg	
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml	
Immunogen	Raw 264 cell line	
External Database Links	UniProt: <u>P30204</u> <u>Related reagents</u> Entrez Gene: <u>20288</u> Msr1 <u>Related reagents</u>	
Synonyms	Scvr	
RRID	AB_323817	
Fusion Partners	Spleen cells from immunized AO rats were fused with cells of	the Y3 rat myeloma cell line
Specificity	Rat anti Mouse CD204 antibody, clone 2F8 recognizes the r class A (SR-A), type I and II, also known as CD204. CD204 is macrophages and functions both as an endocytic receptor for adhesion receptor for macrophages binding to ligand rich tissu lesions. Rat anti Mouse CD204 antibody, clone 2F8 inhibits the low-density lipoproteins and also inhibits divalent cation indepe <i>al.</i> 1993). Rat anti Mouse CD204 antibody, clone 2F8 recognizes an epi polymorphic in the SRA from C57BL/6 mice. Rat anti Mouse CD therefore unsuitable for use with the C57BL/6 mouse strain (D	expressed by tissue lipoproteins and as an ues e.g. atherosclerotic e uptake of acetylated endent adhesion (<u>Fraser <i>et</i></u> itope within SRA that is CD204 antibody, clone 2F8 is
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in	

References

1. Fraser, I. *et al.* (1993) Divalent cation-independent macrophage adhesion inhibited by monoclonal antibody to murine scavenger receptor. <u>Nature. 364 (6435): 343-6.</u>

de Villiers, W.J. *et al.* (1994) Macrophage-colony-stimulating factor selectively enhances macrophage scavenger receptor expression and function. <u>J Exp Med. 180 (2): 705-9.</u>
 Hughes, D.A. *et al.* (1994) Murine M phi scavenger receptor: adhesion function and expression. <u>Immunol Lett. 43 (1-2): 7-14.</u>

4. Hughes, D.A. *et al.* (1995) Murine macrophage scavenger receptor: in vivo expression and function as receptor for macrophage adhesion in lymphoid and non-lymphoid organs. <u>Eur J Immunol. 25 (2): 466-73.</u>

5. Daugherty, A. *et al.* (2000) Polymorphism of class A scavenger receptors in C57BL/6 mice. J Lipid Res. 41 (10): 1568-77.

6. Kaur, H. *et al.* (2003) Identification of a scavenger receptor homologue on nonspecific cytotoxic cells and evidence for binding to oligodeoxyguanosine. <u>Fish Shellfish Immunol.</u> <u>15: 169-81.</u>

7. Kaur, H. *et al.* (2004) Single-base oligodeoxyguanosine-binding proteins on nonspecific cytotoxic cells: identification of a new class of pattern-recognition receptors. <u>Scand J</u> <u>Immunol. 60: 238-48.</u>

8. Allred, K.F. *et al.* (2006) Estrogen receptor-alpha mediates gender differences in atherosclerosis induced by HIV protease inhibitors. <u>J Biol Chem. 281:1419-25.</u>

9. Anatelli, F. *et al.* (2006) Macrophage-targeted photosensitizer conjugate delivered by intratumoral injection. <u>Mol Pharm. 3: 654-64.</u>

10. Luechtenborg, B. *et al.* (2008) Function of scavenger receptor class A type I/II is not important for smooth muscle foam cell formation. <u>Eur J Cell Biol. 87: 91-9.</u>

11. Nielsen, B.S. *et al.* (2008) Matrix metalloproteinase 13 is induced in fibroblasts in polyomavirus middle T antigen-driven mammary carcinoma without influencing tumor progression. <u>PLoS One. 3 (8): e2959.</u>

12. Aid, S. *et al.* (2008) Neuroinflammatory response to lipopolysaccharide is exacerbated in mice genetically deficient in cyclooxygenase-2. J Neuroinflammation. 5: 17.

13. Neyen, C. *et al.* (2009) Macrophage scavenger receptor A mediates adhesion to apolipoproteins A-I and E. <u>Biochemistry. 48 (50): 11858-71.</u>

14. Moldenhauer, L.M. *et al.* (2010) GM-CSF is an essential regulator of T cell activation competence in uterine dendritic cells during early pregnancy in mice. <u>J Immunol. 185 (11):</u> 7085-96.

15. Cainarca, S. *et al.* (2010) A photoprotein in mouse embryonic stem cells measures Ca2+ mobilization in cells and in animals. <u>PLoS One. 5: e8882.</u>

16. Dewals, B.G. *et al.* (2010) IL-4Ralpha-independent expression of mannose receptor and Ym1 by macrophages depends on their IL-10 responsiveness. <u>PLoS Negl Trop Dis.</u> <u>4(5):e689.</u>

17. Swain, S.D. *et al.* (2011) *Pneumocystis* infection in an immunocompetent host can promote collateral sensitization to respiratory antigens. <u>Infect Immun. 79 (5): 1905-14.</u>
18. Hald, A. *et al.* (2011) MMP9 is protective against lethal inflammatory mass lesions in the mouse colon. <u>Dis Model Mech. 4: 212-27.</u>

Nikolic, D. *et al.* (2011) SR-A ligand and M-CSF dynamically regulate SR-A expression and function in primary macrophages via p38 MAPK activation. <u>BMC Immunol. 12: 37.</u>
 Sever-Chroneos, Z. *et al.* (2011) Surfactant Protein A (SP-A)-mediated Clearance of *Staphylococcus aureus* Involves Binding of SP-A to the Staphylococcal Adhesin Eap and the Macrophage Receptors SP-A Receptor 210 and Scavenger Receptor Class A. J Biol

Chem. 286: 4854-70.

21. Zaynagetdinov, R *et al.* (2011) A critical role for macrophages in promotion of urethane-induced lung carcinogenesis. J Immunol. 187 (11): 5703-11.

22. Yang, C.N. *et al.* (2011) Mechanism mediating oligomeric Aβ clearance by naïve primary microglia. <u>Neurobiol Dis. 42 (3): 221-30.</u>

23. Horlad, H. *et al.* (2013) Corosolic acid impairs tumor development and lung metastasis by inhibiting the immunosuppressive activity of myeloid-derived suppressor cells. <u>Mol Nutr</u> Food Res. 57 (6): 1046-54.

24. Koronyo Y *et al.* (2015) Therapeutic effects of glatiramer acetate and grafted CD115+ monocytes in a mouse model of Alzheimer's disease. <u>Brain. 138 (Pt 8): 2399-422.</u>

25. Tao, J. *et al.* (2015) CIC-3 deficiency prevents atherosclerotic lesion development in ApoE^{-/-} mice. J Mol Cell Cardiol. 87: 237-247.

26. Prins, J.R. *et al.* (2015) Unstable Foxp3+ regulatory T cells and altered dendritic cells are associated with lipopolysaccharide-induced fetal loss in pregnant interleukin 10-deficient mice. <u>Biol Reprod. 93 (4): 95.</u>

27. Almholt, K. *et al.* (2015) Spontaneous lung and lymph node metastasis in transgenic breast cancer is independent of the urokinase receptor uPAR. <u>Clin Exp Metastasis. 32 (6):</u> 543-54.

28. Verheijden S *et al.* (2015) Identification of a chronic non-neurodegenerative microglia activation state in a mouse model of peroxisomal β -oxidation deficiency. <u>Glia. 63 (9):</u> <u>1606-20.</u>

29. Tsay, H.J. *et al.* (2016) Identifying N-linked glycan moiety and motifs in the cysteine-rich domain critical for N-glycosylation and intracellular trafficking of SR-AI and MARCO. J Biomed Sci. 23: 27.

30. Fujiwara, Y. *et al.* (2016) Onionin A, a sulfur-containing compound isolated from onions, impairs tumor development and lung metastasis by inhibiting the protumoral and immunosuppressive functions of myeloid cells. <u>Mol Nutr Food Res. 60 (11): 2467-80.</u>

31. Kokubu, Y. *et al.* (2016) Induction of protumoral CD11c(high) macrophages by glioma cancer stem cells through GM-CSF. <u>Genes Cells. 21 (3): 241-51.</u>

32. Sapkota, M. *et al.* (2016) Malondialdehyde-Acetaldehyde-Adducted Surfactant Protein Alters Macrophage Functions Through Scavenger Receptor A. <u>Alcohol Clin Exp Res. 40</u> (12): 2563-2572.

33. Rentsendorj, A. *et al.* (2018) A novel role for osteopontin in macrophage-mediated amyloid- β clearance in Alzheimer's models. <u>Brain Behav Immun. 67: 163-180.</u>

34. Haupt, F. *et al.* (2019) Retinal myeloid cells regulate tip cell selection and vascular branching morphogenesis via Notch ligand Delta-like 1. <u>Sci Rep. 9 (1): 9798.</u>

35. Yamamoto, M. *et al.* (2019) Evaluation of neurobehavioral impairment in methylmercury-treated KK-Ay mice by dynamic weight-bearing test. <u>J Appl Toxicol. 39 (2):</u> 221-30.

36. Cortes, E. *et al.* (2019) GPER is a mechanoregulator of pancreatic stellate cells and the tumor microenvironment. <u>EMBO Rep. 20 (1): e46556.</u>

37. Tian, L.X. *et al.* (2020) Cytochrome P450 1A1 enhances inflammatory responses and impedes phagocytosis of bacteria in macrophages during sepsis. <u>Cell Commun Signal. 18</u> (<u>1): 70</u>.

38. Li, S. *et al.* (2020) Activated Bone Marrow-Derived Macrophages Eradicate
Alzheimer's-Related Aβ(42) Oligomers and Protect Synapses. <u>Front Immunol. 11: 49.</u>
39. Becerril-García, M.A. *et al.* (2020) Langerhans Cells From Mice at Birth Express

	 Endocytic- and Pattern Recognition-Receptors, Migrate to Drain Antigen and Activate Neonatal T Cells in vivo. <u>Front Immunol. 1</u> 40. Koronyo-Hamaoui, M. <i>et al.</i> (2020) Peripherally derived angi enzyme-enhanced macrophages alleviate Alzheimer-related disc <u>336-58.</u> 41. Shiau, D.J. <i>et al.</i> (2020) Hepatocellular carcinoma-derived h triggers M2 macrophage polarization via a TLR2/NOX2/autopha <u>13582.</u> 42. Wang, Y. <i>et al.</i> (2021) Scavenger receptor A1 participates in <i>interrogans</i> serovar Autumnalis strain 56606v and inflammation <u>Emerg Microbes Infect.</u> : 1-39. 43. Mañucat-Tan, N. <i>et al.</i> (2021) Hypochlorite-induced aggrega a novel antioxidant role in blood plasma. <u>Redox Biol. 40</u>: 101847. 44. Jia, J. <i>et al.</i> (2022) Ferritin triggers neutrophil extracellular tr storm through Msr1 contributing to adult-onset Still's disease par <u>13 (1): 6804.</u> 45. Zhang, H. <i>et al.</i> (2024) Ferritin-mediated neutrophil extracell cytokine storm via macrophage scavenger receptor in sepsis-as <u>Commun Signal. 22 (1): 97.</u> 46. Zuroff, L.R. <i>et al.</i> (2020) Effects of IL-34 on Macrophage Imr Response to Alzheimer's-Related Aβ(42) Assemblies. <u>Front Imm</u> 47. Qian, J. <i>et al.</i> (2023) SRA inhibition improves antitumor pote chaperone vaccine. <u>Front Immunol. 14</u>: 1118781. 	1: 744. iotensin converting ease. <u>Brain. 143 (1):</u> igh mobility group box 1 gy axis. <u>Sci Rep. 10 (1):</u> uptake of <i>Leptospira</i> in mouse macrophages. tion of fibrinogen underlies 7. ap-mediated cytokine thogenesis. <u>Nat Commun.</u> ular traps formation and sociated lung injury. <u>Cell</u> munological Profile in <u>nunol. 11: 1449.</u>
Further Reading	1. Gordon, S. (1995) The macrophage. <u>Bioessays. 17 (11): 977-</u>	<u>86.</u>
Storage	Store at -20°C only. This product should be stored undiluted. Storage in frost free freezers is not recommended. Avoid repeat as this may denature the antibody. Should this product contain a recommend microcentrifugation before use.	• •
Guarantee	12 months from date of despatch	
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/MCA1322EL 10162	
Regulatory	For research purposes only	

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16)	DyLight®800
Rabbit Anti Rat IgG (STAR17)	<u>FITC</u>
Goat Anti Rat IgG (STAR73)	<u>RPE</u>
Rabbit Anti Rat IgG (STAR21)	<u>HRP</u>

Goat Ant	i Rat IgG (MOUSE ADS	ORBED) (S ⁻	TAR71) <u>DyLight®550</u> , <u>I</u>	DyLight®650), DyLight®800
Goat Ant	i Rat IgG (STAR131)		<u>Alk. Phos., Bio</u>	<u>tin</u>	
Goat Ant	i Rat IgG (STAR72)		HRP		
Goat Ant	i Rat IgG (STAR69)		<u>FITC</u>		
North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751 Email: antibody_sales_us@bio-ra	id.com	Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-	rad.com	Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M437942:250320'					
Printed on 31 Mar 2025					

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint