

Datasheet: MCA1334GA

BATCH NUMBER 1609

Description:	MOUSE ANTI RAT CD31
Specificity:	CD31
Other names:	PECAM-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	TLD-3A12
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/100
Immunohistology - Frozen	▪			1/10 - 1/100
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting	▪			
Immunofluorescence	▪			
Functional Assays (1)			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) **Bio-Rad recommend the use of [MCA1334EL](#) for use in functional studies**

Target Species

Rat

Species Cross Reactivity

Reacts with: Rhesus Monkey, Pig

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Activated, Lewis rat derived microglial cells.
External Database Links	<p>UniProt: Q3SWT0 Related reagents</p> <p>Entrez Gene: 29583 Pecam1 Related reagents</p>
Synonyms	Pecam
RRID	AB_2161159
Fusion Partners	Spleen cells from immunised BALB/c mouse were fused with cells of the mouse SP2 myeloma cell line.
Specificity	<p>Mouse anti Rat CD31 antibody, clone TLD-3A12 recognizes rat PECAM-1 (CD31), a 661 amino acid type 1 transmembrane protein expressed primarily on endothelial cells, platelets and leucocytes.</p> <p>Clone TLD-3A12 has been shown to partially block the proliferative response of antigen-specific CD4+ T cells to antigen-presenting cells and relevant antigen (Stevenson, K.S. et al.2009).</p> <p>Mouse anti Rat CD31 antibody, clone TLD-3A12 is suitable for use in IHC on formalin-fixed paraffin-embedded sections pre-treated with 0.2M boric acid, pH7.0. (Wilson et al. 2007). Mouse anti Rat CD31, clone TLD-3A12 has been shown to be cross-reactive with endothelial cells derived from rhesus macaque (Maclean et al. 2001)</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	1. Williams, K.C. <i>et al.</i> (1996) PECAM-1 (CD31) expression in the central nervous system and its role in experimental allergic encephalomyelitis in the rat. J Neurosci Res. 45 (6): 747-57.

2. Nakao, A. *et al.* (2003) Carbon monoxide inhalation protects rat intestinal grafts from ischemia/reperfusion injury. [Am J Pathol. 163: 1587-98.](#)
3. Stevenson, K.S. *et al.* (2009) Isolation, characterization, and differentiation of thy1.1-sorted pancreatic adult progenitor cell populations. [Stem Cells Dev. 18 \(10\): 1389-98.](#)
4. Ott, I. *et al.* (2005) Endothelial-like cells expanded from CD34+ blood cells improve left ventricular function after experimental myocardial infarction. [FASEB J. 19 \(8\): 992-4.](#)
5. Fujimoto, K.L. *et al.* (2007) An elastic, biodegradable cardiac patch induces contractile smooth muscle and improves cardiac remodeling and function in subacute myocardial infarction. [J Am Coll Cardiol. 49: 2292-300.](#)
6. Thebault, P. *et al.* (2010) The C-type lectin-like receptor CLEC-1, expressed by myeloid cells and endothelial cells, is up-regulated by immunoregulatory mediators and moderates T cell activation. [J Immunol. 183: 3099-108.](#)
7. Graham, M.J. *et al.* (1998) *In vivo* distribution and metabolism of a phosphorothioate oligonucleotide within rat liver after intravenous administration. [J Pharmacol Exp Ther. 286: 447-58.](#)
8. Haywood, L. *et al.* (2003) Inflammation and angiogenesis in osteoarthritis. [Arthritis Rheum. 48: 2173-7.](#)
9. Kielian, T. and Hickey, W.F. (2010) Proinflammatory cytokine, chemokine, and cellular adhesion molecule expression during the acute phase of experimental brain abscess development. [Am J Pathol. 157: 647-58.](#)
10. Lochhead, J.J. *et al.* (2010) Oxidative stress increases blood-brain barrier permeability and induces alterations in occludin during hypoxia-reoxygenation. [J Cereb Blood Flow Metab. 30: 1625-36.](#)
11. Arkudas, A. *et al.* (2007) Fibrin gel-immobilized VEGF and bFGF efficiently stimulate angiogenesis in the AV loop model. [Mol Med. 13: 480-7.](#)
12. Nakao, A. *et al.* (2011) *Ex vivo* carbon monoxide delivery inhibits intimal hyperplasia in arterialized vein grafts. [Cardiovasc Res. 89: 457-63.](#)
13. Ohnishi, T. *et al.* (2007) Comparison of endothelial cell proliferation in normal liver and adipose tissue in B6C3F1 mice, F344 rats, and humans. [Toxicol Pathol. 35: 904-9.](#)
14. Schilte, M.N. *et al.* (2009) Long-term intervention with heparins in a rat model of peritoneal dialysis. [Perit Dial Int. 29: 26-35.](#)
15. Seegers, H.C. *et al.* (2003) Enhancement of angiogenesis by endogenous substance P release and neurokinin-1 receptors during neurogenic inflammation. [J Pharmacol Exp Ther. 306: 8-12.](#)
16. Wilson, E. *et al.* (2007) An evaluation of the immunohistochemistry benefits of boric acid antigen retrieval on rat decalcified joint tissues. [J Immunol Methods. 322: 137-42.](#)
17. Willis, C.L. *et al.* (2010) Protein kinase C activation modulates reversible increase in cortical blood-brain barrier permeability and tight junction protein expression during hypoxia and posthypoxic reoxygenation. [J Cereb Blood Flow Metab. 30: 1847-59.](#)
18. Salehi-Had, H. *et al.* (2011) Utilizing targeted gene therapy with nanoparticles binding alpha v beta 3 for imaging and treating choroidal neovascularization. [PLoS One. 6: e18864.](#)
19. MacLean, A.G. *et al.* (2001) Rhesus macaque brain microvessel endothelial cells behave in a manner phenotypically distinct from umbilical vein endothelial cells. [J Neuroimmunol. 118: 223-32.](#)
20. Ceelen, W. *et al.* (2007) Recombinant human erythropoietin alpha modulates the effects of radiotherapy on colorectal cancer microvessels. [Br J Cancer. 96: 692-700.](#)

21. Tung, H.C. *et al.* (2015) The Beneficial Effects of P2X7 Antagonism in Rats with Bile Duct Ligation-induced Cirrhosis. [PLoS One. 10 \(5\): e0124654.](#)
22. Oboshi, M. *et al.* (2015) Temporary dietary iron restriction affects the process of thrombus resolution in a rat model of deep vein thrombosis. [PLoS One. 10 \(5\): e0126611.](#)
23. Wu, S.H. *et al.* (2015) Autologous adipose-derived stem cells attenuate muscular atrophy and protect spinal cord ventral horn motor neurons in an animal model of burn injury. [Cytotherapy. 17 \(8\): 1066-75.](#)
24. Ikutomi, M. *et al.* (2015) Diverse contribution of bone marrow-derived late-outgrowth endothelial progenitor cells to vascular repair under pulmonary arterial hypertension and arterial neointimal formation. [J Mol Cell Cardiol. 86: 121-35.](#)
25. Ferrantelli, E. *et al.* (2016) The dipeptide alanyl-glutamine ameliorates peritoneal fibrosis and attenuates IL-17 dependent pathways during peritoneal dialysis. [Kidney Int. 89 \(3\): 625-35.](#)
26. Lux, M. *et al.* (2016) *In vitro* maturation of large-scale cardiac patches based on a perfusable starter matrix by cyclic mechanical stimulation. [Acta Biomater. 30: 177-87.](#)
27. Kotaro, S. *et al.* (2015) Responses of pulp vasculature after cavity preparation in rat molars [Journal of Oral Biosciences. 57 \(3\): 157-64.](#)
28. Teng, B.T. *et al.* (2011) Protective effect of caspase inhibition on compression-induced muscle damage. [J Physiol. 589: 3349-69.](#)
29. Kakaiy, A. *et al.* (2015) Comparing protective effect of grape seed extract versus atorvastatin on endometriosis in rat model: Evidence for immunohistochemical and biochemical alterations. [Vet Res Forum. 6 \(2\): 101-10.](#)
30. Brandl, A. *et al.* (2014) A novel early precursor cell population from rat bone marrow promotes angiogenesis *in vitro*. [BMC Cell Biol. 15: 12.](#)
31. Sun, C.K. *et al.* (2015) Mixed serum-deprived and normal adipose-derived mesenchymal stem cells against acute lung ischemia-reperfusion injury in rats. [Am J Transl Res. 7 \(2\): 209-31.](#)
32. Matsugami, H. *et al.* (2014) VEGF secretion by adipose tissue-derived regenerative cells is impaired under hyperglycemic conditions via glucose transporter activation and ROS increase. [Biomed Res. 35 \(6\): 397-405.](#)
33. Park; J.R. *et al.* (2016) Effects of Peroxisome Proliferator-Activated Receptor- δ Agonist on Cardiac Healing after Myocardial Infarction. [PLoS One. 11 \(2\): e0148510.](#)
34. Naaijkens BA *et al.* (2015) Acute myocardial infarction does not affect functional characteristics of adipose-derived stem cells in rats, but reduces the number of stem cells in adipose tissue. [Cell Tissue Res. 362 \(3\): 623-32.](#)
35. Lim, S. *et al.* (2017) Attenuation of carotid neointimal formation after direct delivery of a recombinant adenovirus expressing glucagon-like peptide-1 in diabetic rats. [Cardiovasc Res. 113 \(2\): 183-94.](#)
36. Stavenuiter, A.W. *et al.* (2015) Protective Effects of Paricalcitol on Peritoneal Remodeling during Peritoneal Dialysis. [Biomed Res Int. 2015: 468574.](#)
37. Frye, C.A. & Patrick, C.W. Jr (2002) Isolation and culture of rat microvascular endothelial cells. [In Vitro Cell Dev Biol Anim. 38 \(4\): 208-12.](#)
38. Jiang, Y. *et al.* (2015) SOD1 nanozyme salvages ischemic brain by locally protecting cerebral vasculature. [J Control Release. 213: 36-44.](#)
39. Mirzaei, M. *et al.* (2017) Nanosilver particles increase follicular atresia: Correlation with oxidative stress and aromatization. [Environ Toxicol. 32 \(10\): 2244-55.](#)
40. Sønstevold, T. *et al.* (2017) Hyperbaric oxygen treatment did not significantly affect

radiation injury in the mandibular area of rats. [Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology. \[Epub ahead of print\].](#)

41. Ichihara, Y. *et al.* (2018) Self-assembling peptide hydrogel enables instant epicardial coating of the heart with mesenchymal stromal cells for the treatment of heart failure.

[Biomaterials. 154: 12-23.](#)

42. Melly, L. *et al.* (2018) Myocardial infarction stabilization by cell-based expression of controlled Vascular Endothelial Growth Factor levels. [J Cell Mol Med. 22 \(5\): 2580-91.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1334GA>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M352889:190408'

