

Datasheet: MCA1335GA

BATCH NUMBER 166374

Description:	MOUSE ANTI PIG SLA CLASS II DQ
Specificity:	SLA CLASS II DQ
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	K274.3G8
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig
Species Cross Reactivity	<p>Reacts with: Bovine</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Porcine peripheral blood lymphocytes
Fusion Partners	Spleen cells from immunized mice were fused with cells of the P3-X63-Ag.653 myeloma cell line
Specificity	Mouse anti Pig SLA Class II DQ antibody, clone K274.3G8 recognizes SLA DQ molecules which are expressed on all B cells, antigen presenting cells and on certain subsets of resting and activated T cells. The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In pigs, this is referred to as the swine leukocyte antigen (SLA) region. There are 3 major MHC class II proteins encoded by the SLA which are SLA DP, SLA DQ and SLA DR.
Flow Cytometry	Use 10µl of the suggested working dilution to label 1x10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> 1. Brodersen, R. <i>et al.</i> (1998) Analysis of the immunological cross reactivities of 213 well characterized monoclonal antibodies with specificities against various leucocyte surface antigens of human and 11 animal species. Vet Immunol Immunopathol. 64: 1-13. 2. Paillot, R. <i>et al.</i> (2001) Functional and phenotypic characterization of distinct porcine dendritic cells derived from peripheral blood monocytes. Immunology 102: 396-404. 3. Makala, L.H. <i>et al.</i> (2001) Ontogeny of pig discrete Peyer's patches: expression of surface antigens. J Vet Med Sci. 63 (6): 625-36. 4. Faure, J.P. <i>et al.</i> (2002) Polyethylene glycol reduces early and long-term cold ischemia-reperfusion and renal medulla injury. J Pharmacol Exp Ther. 302: 861-70. 5. Hauet, T. <i>et al.</i> (2002) Polyethylene glycol reduces the inflammatory injury due to cold ischemia/reperfusion in autotransplanted pig kidneys. Kidney Int. 62: 654-67. 6. Yang, P. <i>et al.</i> (2002) Immune cells in the porcine retina: distribution, characterization and morphological features. Invest Ophthalmol Vis Sci. 43: 1488-92. 7. Sarradell, J. <i>et al.</i> (2003) A morphologic and immunohistochemical study of the bronchus-associated lymphoid tissue of pigs naturally infected with <i>Mycoplasma hyopneumoniae</i>. Vet Pathol. 40: 395-404. 8. Edamura, K. <i>et al.</i> (2005) Effect of long-term culture on the expression of antigens and adhesion molecule in single porcine pancreatic endocrine cells. Xenotransplantation. 12 (4): 327-32. 9. Maasilta, P.K. <i>et al.</i> (2005) Immune cells in a heterotopic lamb-to-pig bronchial xenograft model. Transpl Int. 18: 1100-8. 10. Jayle, C. <i>et al.</i> (2007) Comparison of protective effects of trimetazidine against experimental warm ischemia of different durations: early and long-term effects in a pig kidney model. Am J Physiol Renal Physiol. 292: F1082-93.

11. Park, J.Y. *et al.* (2008) Characterization of interaction between porcine reproductive and respiratory syndrome virus and porcine dendritic cells. [J Microbiol Biotechnol. 18: 1709-16.](#)
12. Facci, M.R. *et al.* (2010) A comparison between isolated blood dendritic cells and monocyte-derived dendritic cells in pigs. [Immunology. 129 \(3\): 396-405.](#)
13. Inman, C.F. *et al.* (2010) Dendritic cells interact with CD4 T cells in intestinal mucosa. [J Leukoc Biol. 88: 571-8.](#)
14. Weesendorp E *et al.* (2013) Phenotypic modulation and cytokine profiles of antigen presenting cells by European subtype 1 and 3 porcine reproductive and respiratory syndrome virus strains *in vitro* and *in vivo*. [Vet Microbiol. 167 \(3-4\): 638-50.](#)
15. Debeer, S. *et al.* (2013) Comparative histology and immunohistochemistry of porcine versus human skin. [Eur J Dermatol. 23 \(4\): 456-66.](#)
16. LeLuduec, J.B. *et al.* (2016) Intradermal vaccination with un-adjuvanted sub-unit vaccines triggers skin innate immunity and confers protective respiratory immunity in domestic swine. [Vaccine. 34 \(7\): 914-22.](#)
17. Vreman, S. *et al.* (2018) Neonatal porcine blood derived dendritic cell subsets show activation after TLR2 or TLR9 stimulation. [Dev Comp Immunol. 84: 361-70.](#)
18. Loss, H. *et al.* (2018) Effects of a pathogenic ETEC strain and a probiotic *Enterococcus faecium* strain on the inflammasome response in porcine dendritic cells. [Vet Immunol Immunopathol. 203: 78-87.](#)
19. Ladowski, J.M. *et al.* (2018) Swine Leukocyte Antigen Class II Is a Xenoantigen. [Transplantation. 102 \(2\): 249-54.](#)

Further Reading	1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. Vet Res. 39: 54.
Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1335GA 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

- Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
- Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)
- Goat Anti Mouse IgG (STAR76...) [RPE](#)

Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M409228:221018'

Printed on 18 Jan 2024