

## Datasheet: MCA1335GA

### BATCH NUMBER 1808

<b>Description:</b>	MOUSE ANTI PIG SLA CLASS II DQ
<b>Specificity:</b>	SLA CLASS II DQ
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	K274.3G8
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Pig
<b>Species Cross Reactivity</b>	<p>Reacts with: Bovine</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Porcine peripheral blood lymphocytes
<b>Fusion Partners</b>	Spleen cells from immunized mice were fused with cells of the P3-X63-Ag.653 myeloma cell line
<b>Specificity</b>	<b>Mouse anti Pig SLA Class II DQ antibody, clone K274.3G8</b> recognizes SLA DQ molecules which are expressed on all B cells, antigen presenting cells and on certain subsets of resting and activated T cells. The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In pigs, this is referred to as the swine leukocyte antigen (SLA) region. There are 3 major MHC class II proteins encoded by the SLA which are SLA DP, SLA DQ and SLA DR.
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul
<b>References</b>	<ol style="list-style-type: none"> <li>1. Lunney, J.K. (1993) Characterization of swine leukocyte differentiation antigens. <a href="#">Immunol Today 14: 147-8.</a></li> <li>2. Brodersen, R. <i>et al.</i> (1998) Analysis of the immunological cross reactivities of 213 well characterized monoclonal antibodies with specificities against various leucocyte surface antigens of human and 11 animal species. <a href="#">Vet Immunol Immunopathol. 64: 1-13.</a></li> <li>3. Sarradell, J. <i>et al.</i> (2003) A morphologic and immunohistochemical study of the bronchus-associated lymphoid tissue of pigs naturally infected with <i>Mycoplasma hyopneumoniae</i>. <a href="#">Vet Pathol. 40: 395-404.</a></li> <li>4. Inman, C.F. <i>et al.</i> (2010) Dendritic cells interact with CD4 T cells in intestinal mucosa. <a href="#">J Leukoc Biol. 88: 571-8.</a></li> <li>5. Faure, J.P. <i>et al.</i> (2002) Polyethylene glycol reduces early and long-term cold ischemia-reperfusion and renal medulla injury. <a href="#">J Pharmacol Exp Ther. 302: 861-70.</a></li> <li>6. Hauet, T. <i>et al.</i> (2002) Polyethylene glycol reduces the inflammatory injury due to cold ischemia/reperfusion in autotransplanted pig kidneys. <a href="#">Kidney Int. 62: 654-67.</a></li> <li>7. Paillot, R. <i>et al.</i> (2001) Functional and phenotypic characterization of distinct porcine dendritic cells derived from peripheral blood monocytes. <a href="#">Immunology 102: 396-404.</a></li> <li>8. Yang, P. <i>et al.</i> (2002) Immune cells in the porcine retina: distribution, characterization and morphological features. <a href="#">Invest Ophthalmol Vis Sci. 43: 1488-92.</a></li> <li>9. Jayle, C. <i>et al.</i> (2007) Comparison of protective effects of trimetazidine against experimental warm ischemia of different durations: early and long-term effects in a pig kidney model. <a href="#">Am J Physiol Renal Physiol. 292: F1082-93.</a></li> <li>10. Park, J.Y. <i>et al.</i> (2008) Characterization of interaction between porcine reproductive and respiratory syndrome virus and porcine dendritic cells. <a href="#">J Microbiol Biotechnol. 18: 1709-16.</a></li> </ol>

11. Maasilta, P.K. *et al.* (2005) Immune cells in a heterotopic lamb-to-pig bronchial xenograft model. [Transpl Int. 18: 1100-8.](#)
12. Weesendorp E *et al.* (2013) Phenotypic modulation and cytokine profiles of antigen presenting cells by European subtype 1 and 3 porcine reproductive and respiratory syndrome virus strains *in vitro* and *in vivo*. [Vet Microbiol. 167 \(3-4\): 638-50.](#)
13. Makala, L.H. *et al.* (2001) Ontogeny of pig discrete Peyer's patches: expression of surface antigens. [J Vet Med Sci. 63 \(6\): 625-36.](#)
14. Facci, M.R. *et al.* (2010) A comparison between isolated blood dendritic cells and monocyte-derived dendritic cells in pigs. [Immunology. 129 \(3\): 396-405.](#)
15. Edamura, K. *et al.* (2005) Effect of long-term culture on the expression of antigens and adhesion molecule in single porcine pancreatic endocrine cells. [Xenotransplantation. 12 \(4\): 327-32.](#)
16. Debeer, S. *et al.* (2013) Comparative histology and immunohistochemistry of porcine versus human skin. [Eur J Dermatol. 23 \(4\): 456-66.](#)
17. Loss, H. *et al.* (2018) Effects of a pathogenic ETEC strain and a probiotic *Enterococcus faecium* strain on the inflammasome response in porcine dendritic cells. [Vet Immunol Immunopathol. 203: 78-87.](#)
18. Vreman, S. *et al.* (2018) Neonatal porcine blood derived dendritic cell subsets show activation after TLR2 or TLR9 stimulation. [Dev Comp Immunol. 84: 361-70.](#)
19. LeLuduec, J.B. *et al.* (2016) Intradermal vaccination with un-adjuvanted sub-unit vaccines triggers skin innate immunity and confers protective respiratory immunity in domestic swine. [Vaccine. 34 \(7\): 914-22.](#)

---

**Further Reading**      1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)

---

**Storage**                      Store at +4°C or at -20°C if preferred.

   This product should be stored undiluted.

   Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**                      12 months from date of despatch

---

**Health And Safety Information**      Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1335GA>  
10040

---

**Regulatory**                      For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

- |                                   |                     |
|-----------------------------------|---------------------|
| Goat Anti Mouse IgG (STAR77...)   | <a href="#">HRP</a> |
| Rabbit Anti Mouse IgG (STAR12...) | <a href="#">RPE</a> |

Goat Anti Mouse IgG (STAR70...) [FITC](#)  
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)  
Goat Anti Mouse IgG (STAR76...) [RPE](#)  
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),  
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),  
[FITC](#), [HRP](#)  
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)  
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)  
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

## Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M365171:200529'

Printed on 25 Mar 2023

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)