

Datasheet: MCA1340F

Description:	MOUSE ANTI HUMAN CD120a:FITC
Specificity:	CD120a
Other names:	TNF-R1
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	H398
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human			
Species Cross	Reacts with: Rabb	pit		
Reactivity	N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications of personal communications from the originators. Please refer to references indicated further information.			viewed publications or
Product Form	Purified IgG conju	gated to Fluorescein Isoth	niocyanate Isomer 1	(FITC) - liquid
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
	FITC	490	525	
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant			
Buffer Solution	Phosphate buffere			

Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	Purified human tumour necrosis factor receptor type 1.		
External Database			
Links	UniProt: P19438 Related reagents		
	Entrez Gene:		
	7132 TNFRSF1A Related reagents		
Synonyms	TNFAR, TNFR1		
RRID	AB_322825		
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS0 myeloma cell line.		
Specificity	Mouse anti Human CD120a antibody, clone H398 recognizes an extracellular domain of the ~55 kDa human TNF receptor (p55, TNF-R1, CD120a). No binding occurs to the ~75 kDa TNF receptor (CD120b). CD120a is weakly expressed by monocytes and granulocytes.		
	Mouse anti Human CD120a antibody, clone H398 may be used to detect high levels of TNFR1, in western blotting under reducing conditions, such as recombinant material, but it is not suitable for detection of TNFR1 in normal cells in this application.		
	Mouse anti Human CD120a antibody, clone H398 inhibits the biological activity of both natural and recombinant human TNFalpha and TNFbeta (<u>Thoma et al. 1990</u> and <u>Dri et al. 1999</u>).		
	Mouse anti Human CD120a antibody, clone H398 is routinely tested in flow cytometry on human peripheral blood monocytes.		
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood.		
References	1. Thoma, B. <i>et al.</i> (1990) Identification of a 60-kD tumor necrosis factor (TNF) receptor as the major signal transducing component in TNF responses. <u>J Exp Med. 172 (4): 1019-23.</u> 2. Menegazzi, R. <i>et al.</i> (1994) Evidence that tumor necrosis factor alpha (TNF)-induced activation of neutrophil respiratory burst on biologic surfaces is mediated by the p55 TNF receptor. <u>Blood. 84 (1): 287-93.</u>		

4. Kohrgruber, N. et al. (1999) Survival, maturation, and function of CD11c- and CD11c+

neutrophil respiratory burst. <u>J Immunol. 162 (1): 460-6.</u>

peripheral blood dendritic cells are differentially regulated by cytokines. <u>J Immunol. 163</u> (6): 3250-9.

- 5. Weigert, N. *et al.* (1996) Gastrin secretion from primary cultures of rabbit antral G cells: stimulation by inflammatory cytokines. <u>Gastroenterology</u>. 110 (1): 147-54.
- 6. Kennedy, G. *et al.* (2010) Biochemical and vascular aspects of pediatric chronic fatigue syndrome. Arch Pediatr Adolesc Med. 164 (9): 817-23.
- 7. Gregory, A.P. *et al.* (2012) TNF receptor 1 genetic risk mirrors outcome of anti-TNF therapy in multiple sclerosis. <u>Nature</u>. 488 (7412): 508-11.
- 8. Schett, G. *et al.* (2003) TNFalpha mediates susceptibility to heat-induced apoptosis by protein phosphatase-mediated inhibition of the HSF1/hsp70 stress response. <u>Cell Death Differ.</u> 10: 1126-36.
- 9. Buckley, C.D. *et al.* (2005) Identification of a phenotypically and functionally distinct population of long-lived neutrophils in a model of reverse endothelial migration. <u>J Leukoc Biol.79</u>: 303-11.
- 10. Thiery, J. *et al.* (2003) Potentiation of a tumor cell susceptibility to autologous CTL killing by restoration of wild-type p53 function. J Immunol. 170: 5919-26.
- 11. Lin, K.H. *et al.* (2009) Mechanisms of resveratrol-induced platelet apoptosis. Cardiovasc Res. 83: 575-85.
- 12. Baker, P.K. *et al.* (2012) Response of hairy cells to IFN-alpha involves induction of apoptosis through autocrine TNF-alpha and protection by adhesion. <u>Blood. 100: 647-53.</u>
- 13. Nakamura-Lopez, Y. *et al.* (2015) RSV P-protein impairs extrinsic apoptosis pathway in a macrophage-like cell line persistently infected with respiratory syncytial virus. <u>Virus Res. 204: 82-7.</u>

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL:FITC (MCA929F)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)
 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739 Fax: +49 (0) 89 8090 95 50 From Email: antibody_sales_us@bio-rad.com Email: antibody_sales_uk@bio-rad.com Email: antibody_sales_de@bio-rad.com

15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M372610:200708'

Printed on 09 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint