Datasheet: MCA1340T BATCH NUMBER 160120

Description:	MOUSE ANTI HUMAN CD120a		
Specificity:	CD120a		
Other names:	TNF-R1		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	H398		
lsotype:	lgG2a		
Quantity:	25 µg		

Product Details

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen			•	
Immunohistology - Paraffin				
ELISA				
Immunoprecipitation			•	
Western Blotting				
Radioimmunoassays				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Reacts with: Rabbit N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture

	supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml		
Immunogen	Purified human tumour necrosis factor receptor type 1.		
External Database Links	UniProt: <u>P19438</u> <u>Related reagents</u> Entrez Gene: <u>7132</u> TNFRSF1A <u>Related reagents</u>		
Synonyms	TNFAR, TNFR1		
RRID	AB_2271880		
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS0 myeloma cell line.		
Specificity	Mouse anti Human CD120a antibody, clone H398 recognizes an extracellular domain of the ~55 kDa human TNF receptor (p55, TNF-R1, CD120a). No binding occurs to the ~75 kDa TNF receptor (CD120b). CD120a is weakly expressed by monocytes and granulocytes.		
Specificity	the ~55 kDa human TNF receptor (p55, TNF-R1, CD120a). No binding occurs to the ~75 kDa TNF receptor (CD120b). CD120a is weakly expressed by monocytes and		
Specificity	 the ~55 kDa human TNF receptor (p55, TNF-R1, CD120a). No binding occurs to the ~75 kDa TNF receptor (CD120b). CD120a is weakly expressed by monocytes and granulocytes. Mouse anti Human CD120a antibody, clone H398 may be used to detect high levels of TNFR1, in western blotting under reducing conditions, such as recombinant material, but it 		
Specificity	 the ~55 kDa human TNF receptor (p55, TNF-R1, CD120a). No binding occurs to the ~75 kDa TNF receptor (CD120b). CD120a is weakly expressed by monocytes and granulocytes. Mouse anti Human CD120a antibody, clone H398 may be used to detect high levels of TNFR1, in western blotting under reducing conditions, such as recombinant material, but it is not suitable for detection of TNFR1 in normal cells in this application. Mouse anti Human CD120a antibody, clone H398 inhibits the biological activity of both natural and recombinant human TNFalpha and TNFbeta (Thoma <i>et al.</i> 1990 and Dri <i>et al.</i> 		

	 4. Kohrgruber, N. <i>et al.</i> (1999) Survival, maturation, and function of CD11c- and CD11c+ peripheral blood dendritic cells are differentially regulated by cytokines. J Immunol. 163 (6): 3250-9. 5. Weigert, N. <i>et al.</i> (1996) Gastrin secretion from primary cultures of rabbit antral G cells: stimulation by inflammatory cytokines. Gastroenterology. 110 (1): 147-54. 6. Kennedy, G. <i>et al.</i> (2010) Biochemical and vascular aspects of pediatric chronic fatigue syndrome. Arch Pediatr Adolesc Med. 164 (9): 817-23. 7. Gregory, A.P. <i>et al.</i> (2012) TNF receptor 1 genetic risk mirrors outcome of anti-TNF therapy in multiple sclerosis. Nature. 488 (7412): 508-11. 8. Schett, G. <i>et al.</i> (2003) TNFalpha mediates susceptibility to heat-induced apoptosis by protein phosphatase-mediated inhibition of the HSF1/hsp70 stress response. Cell Death Differ. 10: 1126-36. 9. Buckley, C.D. <i>et al.</i> (2005) Identification of a phenotypically and functionally distinct population of long-lived neutrophils in a model of reverse endothelial migration. J Leukoc Biol.79: 303-11. 10. Thiery, J. <i>et al.</i> (2003) Potentiation of a tumor cell susceptibility to autologous CTL killing by restoration of wild-type p53 function. J Immunol. 170: 5919-26. 11. Lin, K.H. <i>et al.</i> (2012) Response of hairy cells to IFN-alpha involves induction of apoptosis through autocrine TNF-alpha and protection by adhesion. Blood. 100: 647-53. 13. Nakamura-Lopez, Y. <i>et al.</i> (2015) RSV P-protein impairs extrinsic apoptosis pathway in a macrophage-like cell line persistently infected with respiratory syncytial virus. <u>Virus Res. 204: 82-7.</u> 14. Yang, F. <i>et al.</i> (2013) Improved retinal ganglion cell survival through retinal microglia suppression by a chinese herb extract, triptolide, in the DBA/2J mouse model of glaucoma. <u>Ocul Immunol 1nflamm. 21 (5): 378-89.</u>
Storage	Store at +4°C or at -20°C if preferred.
	This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1340T 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)RPEGoat Anti Mouse IgG IgA IgM (STAR87...)HRP

Goat Anti Mouse IgG (STAR7	6) RP	ΡE			
Goat Anti Mouse IgG (STAR70)					
Rabbit Anti Mouse IgG (STAR13)		<u>RP</u>			
Goat Anti Mouse IgG (Fc) (STAR120)		FITC, HRP			
Rabbit Anti Mouse IgG (STAR9)		<u>rc</u>			
Goat Anti Mouse IgG (STAR7	7) <u>HR</u>	<u>RP</u>			
Goat Anti Mouse IgG (H/L) (S	TAR117) <u>Alk</u>	<u>. Phos., DyLight®488, I</u>	DyLight®550	,	
	Dy	Light®650, DyLight®68	0, <u>DyLight®8</u>	<u>00</u> ,	
	<u>F1</u>	<u>C, HRP</u>			
North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751 Email: antibody_sales_us@l	Worldwide bio-rad.com	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio	Europe -rad.com	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com	
To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M372367:200702'					
Printed on 20 Mar 2025					

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint