

## Datasheet: MCA1352

<b>Description:</b>	MOUSE ANTI GST
<b>Specificity:</b>	GST
<b>Other names:</b>	GLUTATHIONE-S-TRANSFERASE
<b>Format:</b>	S/N
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	vpg66
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	2 ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			1/100 - 1/500
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Product Form</b>	Tissue Culture Supernatant - liquid
<b>Preparation</b>	Tissue Culture Supernatant containing 10mM Tris/HCl pH7.5 and 5-10% foetal calf serum
<b>Preservative Stabilisers</b>	0.1% Sodium Azide
<b>Immunogen</b>	GST from <i>Schistosoma japonicum</i> .
<b>RRID</b>	AB_322081
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS0

myeloma cell line.

---

<b>Specificity</b>	<b>Mouse anti GST antibody, clone vpg66</b> reacts with Glutathione-S-transferase
--------------------	---

---

<b>References</b>	<ol style="list-style-type: none"><li>1. Harris, M.P. &amp; Neil, J.C. (1994) Myristoylation-dependent binding of HIV-1 Nef to CD4. <a href="#">J Mol Biol. 241 (2): 136-42.</a></li><li>2. Blüggel, M. <i>et al.</i> (2011) Toward protein biomarkers for allergy: CD4+ T cell proteomics in allergic and nonallergic subjects sampled in and out of pollen season. <a href="#">J Proteome Res. 10 (4): 1558-70.</a></li><li>3. Shelton, H. &amp; Harris, M. (2008) Hepatitis C virus NS5A protein binds the SH3 domain of the Fyn tyrosine kinase with high affinity: mutagenic analysis of residues within the SH3 domain that contribute to the interaction. <a href="#">Virology 378: 24-32.</a></li><li>4. Foster, T.L. <i>et al.</i> (2011) Cyclophilin A interacts with domain II of hepatitis C virus NS5A and stimulates RNA binding in an isomerase-dependent manner. <a href="#">J Virol. 85: 7460-4.</a></li><li>5. Elfgang, C. <i>et al.</i> (1999) Evidence for specific nucleocytoplasmic transport pathways used by leucine-rich nuclear export signals. <a href="#">Proc Natl Acad Sci U S A. 96: 6229-34.</a></li><li>6. Davis, M.P. (2008) Recombinant VP4 of human rhinovirus induces permeability in model membranes. <a href="#">J Virol. 82: 4169-74.</a></li><li>7. Heger, P. <i>et al.</i> (1998) Multimer formation is not essential for nuclear export of human T-cell leukemia virus type 1 Rex trans-activator protein. <a href="#">J Virol. 72: 8659-68.</a></li><li>8. Lischka, P. <i>et al.</i> (2001) A novel transferable nuclear export signal mediates CRM1-independent nucleocytoplasmic shuttling of the human cytomegalovirus transactivator protein pUL69. <a href="#">EMBO J. 20: 7271-83.</a></li></ol>
-------------------	--

---

<b>Storage</b>	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>
----------------	---

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10336 available at: 10336: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10336.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10336.pdf</a>
--------------------------------------	---

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR8...)	<a href="#">DyLight®800</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>

Rabbit Anti Mouse IgG (STAR13...) [HRP](#)  
Goat Anti Mouse IgG (STAR70...) [FITC](#)  
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)  
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),  
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),  
[FITC](#), [HRP](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M387972:210803'

Printed on 06 Jan 2022

---

© 2022 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)