

Datasheet: MCA1352B

Description:	MOUSE ANTI GST:Biotin
Specificity:	GST
Other names:	GLUTATHIONE-S-TRANSFERASE
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	vpg66
Isotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications	This product has been re	eported to	work in th	e following applications	s. This information is
	derived from testing with	in our lab	oratories, p	peer-reviewed publication	ions or personal
	communications from the	e originato	rs. Please	refer to references inc	dicated for further
	information. For general	protocol r	ecommend	dations, please visit <u>wv</u>	<u>vw.bio-</u>
	rad-antibodies.com/proto	ocols.			
		Voc	No	Not Determined	Suggested Dilution

		0010.				
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry					
	Immunohistology - Frozen					
	Immunohistology - Paraffin			•		
	ELISA	-				
	Immunoprecipitation					
	Western Blotting	•			1/100 - 1/500	
	Where this product has r	not been t	ested for	use in a particular tech	nnique this does not	
	necessarily exclude its us					
	•		•		• •	
	a guide only. It is recommended that the user titrates the product for use in their own					
	system using appropriate	e negative	positive (controls.		
Product Form	Purified IgG conjugated t	o biotin -	liquid			
Preparation	Antibody purified from tis	sue cultu	re supern	atant		
Buffer Solution	Phosphate buffered salin	e				
Preservative Stabilisers	0.09% sodium azide (Na	N ₃)				
Approx. Protein Concentrations	IgG concentration 0.1mg	/ml				
Immunogen	GST from <i>Schistosoma j</i>	aponicum	l.			

RRID		AB_322967				
Fusion P	Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS0 myeloma cell line.				
Specifici	ty	Mouse anti GST antibody, clone vpg66 reacts with Glutathione-S-transferase				
Referenc	es	 Harris, M.P. & Neil, J.C. (1994) Myristoylation-dependent binding of HIV-1 Nef to CD4 Mol Biol. 241 (2): 136-42. Heger, P. <i>et al.</i> (1998) Multimer formation is not essential for nuclear export of humar T-cell leukemia virus type 1 Rex trans-activator protein. J Virol. 72: 8659-68. Elfgang, C. <i>et al.</i> (1999) Evidence for specific nucleocytoplasmic transport pathways used by leucine-rich nuclear export signals. Proc Natl Acad Sci U S A. 96: 6229-34. Lischka, P. <i>et al.</i> (2001) A novel transferable nuclear export signal mediates CRM1-independent nucleocytoplasmic shuttling of the human cytomegalovirus transactivator protein pUL69. EMBO J. 20: 7271-83. Davis, M.P. (2008) Recombinant VP4 of human rhinovirus induces permeability in model membranes. J Virol. 82: 4169-74. Shelton, H. & Harris, M. (2008) Hepatitis C virus NS5A protein binds the SH3 domain the Fyn tyrosine kinase with high affinity: mutagenic analysis of residues within the SH3 domain that contribute to the interaction. <u>Virol J. 5: 24.</u> Foster, T.L. <i>et al.</i> (2011) Cyclophilin A interacts with domain II of hepatitis C virus NS5 and stimulates RNA binding in an isomerase-dependent manner. J Virol. 85: 7460-4. Blüggel, M. <i>et al.</i> (2011) Toward protein biomarkers for allergy: CD4+ T cell proteomic in allergic and nonallergic subjects sampled in and out of pollen season. J Proteome Ref 10 (4): 1558-70. 				
Storage		This product is shipped at ambient temperature. It is recommended to aliquot and store -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C t short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in				
		frost-free freezers is not recommended.				
Guarante	9e	12 months from date of despatch				
	nd Safety ion	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1352B 10040				
Informati						
Informati Regulato	ory	For research purposes only				

'M410376:221028'

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint