

Datasheet: MCA1365G BATCH NUMBER 171593

| Description: | RAT ANTI MOUSE CD117 | |
|---------------|----------------------|--|
| Specificity: | CD117 | |
| Other names: | C-KIT, SCF-R | |
| Format: | Purified | |
| Product Type: | Monoclonal Antibody | |
| Clone: | 2B8 | |
| Isotype: | lgG2b | |
| Quantity: | 0.25 mg | |
| | | |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | | | | 1/50 - 1/200 |
| Immunohistology - Frozen | | | | |
| Immunohistology - Paraffin | | | • | |
| ELISA | | | | |
| Immunoprecipitation | | | | |
| Western Blotting | | | • | |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| Target Species | Mouse | |
|-----------------------------|--|---------------------|
| Product Form | Purified IgG - liquid | |
| Preparation | Purified IgG prepared by affinity chromatography on Protein G supernatant | From tissue culture |
| Buffer Solution | Phosphate buffered saline | |
| Preservative Stabilisers | 0.09% sodium azide (NaN ₃) | |

| Carrier Free | Yes | | | |
|-----------------------------------|---|--|--|--|
| Approx. Protein Concentrations | IgG concentration 1.0 mg/ml | | | |
| Immunogen | IL-3 dependent mast cells. | | | |
| External Database Links | UniProt: P05532 Related reagents Entrez Gene: 16590 Kit Related reagents | | | |
| Synonyms | SI | | | |
| RRID | AB_844471 | | | |
| Fusion Partners | Spleen cells from immunised Wistar rats were fused with cells of the Fox-NY mouse myeloma cell line. | | | |
| Specificity | Rat anti Mouse CD117 antibody, clone 2B8 recognizes mouse CD117, a ~150 kDa cell surface molecule otherwise known as the Kit, or c-kit, receptor. Rat anti Mouse CD117 antibody, clone 2B8 recognizes an epitope within the extracellular region of CD117. In mice, CD117 is predominantly expressed on multipotent haematopoietic stem cells from the adult bone marrow and foetal liver. | | | |
| Flow Cytometry | Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl | | | |
| Further Reading | 1. Manova, K. <i>et al.</i> (1992) c-kit receptor and ligand expression in postnatal development of the mouse cerebellum suggests a function for c-kit in inhibitory interneurons. <u>J Neurosci. 12 (12): 4663-76.</u> 2. Maeda, H. <i>et al.</i> (1992) Requirement of c-kit for development of intestinal pacemaker system. <u>Development. 116 (2): 369-75.</u> 3. Takagi, M. <i>et al.</i> (1992) Stimulation of mouse connective tissue-type mast cells by hemopoietic stem cell factor, a ligand for the c-kit receptor. <u>J Immunol. 148 (11): 3446-53.</u> 4. de Vries, P. <i>et al.</i> (1992) Thymus reconstitution by c-kit-expressing hematopoietic stem cells purified from adult mouse bone marrow. <u>J Exp Med. 176 (6): 1503-9.</u> 5. Fleming WH <i>et al.</i> (1993) Steel factor influences the distribution and activity of murine hematopoietic stem cells <i>in vivo</i> . <u>Proc Natl Acad Sci U S A. 90 (8): 3760-4.</u> 6. Rico-Vargas, S.A. <i>et al.</i> (1994) c-kit expression by B cell precursors in mouse bone marrow. Stimulation of B cell genesis by in vivo treatment with anti-c-kit antibody. <u>J Immunol. 152 (6): 2845-52.</u> | | | |
| Storage | This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. | | | |

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

| Guarantee | 12 months from date of despatch |
|----------------------------------|--|
| Health And Safety Information | Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1365G |
| Regulatory | For research purposes only |

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...) <u>DyLight®800</u>

Rabbit Anti Rat IgG (STAR17...) FITC

Goat Anti Rat IgG (STAR73...) RPE

Rabbit Anti Rat IgG (STAR21...) HRP

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) <u>DyLight®550</u>, <u>DyLight®650</u>, <u>DyLight®800</u>

Goat Anti Rat IgG (STAR131...) Alk. Phos., Biotin

Goat Anti Rat IgG (STAR72...)

HRP

Goat Anti Rat IgG (STAR69...)

FITC

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M409371:221018'

Printed on 03 Sep 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint