

Datasheet: MCA1365SBUV605

Description:	RAT ANTI MOUSE CD117:StarBright UltraViolet 605
Specificity:	CD117
Other names:	C-KIT, SCF-R
Format:	StarBright UltraViolet 605
Product Type:	Monoclonal Antibody
Clone:	2B8
Isotype:	lgG2b
Quantity:	100 TESTS/0.5ml

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry	-			Neat		
	Where this product has	s not been test	ed for u	use in a particular tech	inique this does not		
	necessarily exclude its a guide only. It is recor system using appropria	mmended that	the use	er titrates the product f	g dilutions are given as or use in their own		
Target Species	Mouse						
Product Form	Purified IgG conjugate	d to StarBright	UltraVi	olet 605 - liquid			
Max Ex/Em	Fluorophore	Excitation Max	x (nm)	Emission Max (nm)			
	StarBright UltraViolet 605	340		609			
Preparation	Purified IgG prepared supernatant	by affinity chro	matogra	aphy on Protein G fror	n tissue culture		
Buffer Solution	Phosphate buffered sa	line					
Preservative Stabilisers	0.09% sodium azide (1 1% bovine serum albu 0.1% Pluronic F68 0.1% PEG 3350	0,					

	0.05% Tween 20
Immunogen	IL-3 dependent mast cells.
External Database Links	UniProt: P05532 Related reagents Entrez Gene: 16590 Kit Related reagents
Synonyms	SI
Fusion Partners	Spleen cells from immunised Wistar rats were fused with cells of the Fox-NY mouse myeloma cell line.
Specificity	Rat anti Mouse CD117 antibody, clone 2B8 recognizes mouse CD117, a ~150 kDa cell surface molecule otherwise known as the Kit, or c-kit, receptor.
	Rat anti Mouse CD117 antibody, clone 2B8 recognizes an epitope within the extracellular region of CD117. In mice, CD117 is predominantly expressed on multipotent haematopoietic stem cells from the adult bone marrow and foetal liver.
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 ⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
Further Reading	 Manova, K. <i>et al.</i> (1992) c-kit receptor and ligand expression in postnatal development of the mouse cerebellum suggests a function for c-kit in inhibitory interneurons. J <u>Neurosci. 12 (12): 4663-76.</u> Maeda, H. <i>et al.</i> (1992) Requirement of c-kit for development of intestinal pacemaker system. <u>Development. 116 (2): 369-75.</u> Takagi, M. <i>et al.</i> (1992) Stimulation of mouse connective tissue-type mast cells by hemopoietic stem cell factor, a ligand for the c-kit receptor. <u>J Immunol. 148 (11): 3446-53.</u> de Vries, P. <i>et al.</i> (1992) Thymus reconstitution by c-kit-expressing hematopoietic stem cells purified from adult mouse bone marrow. <u>J Exp Med. 176 (6): 1503-9.</u> Fleming WH <i>et al.</i> (1993) Steel factor influences the distribution and activity of murine hematopoietic stem cells <i>in vivo</i>. <u>Proc Natl Acad Sci U S A. 90 (8): 3760-4.</u> Rico-Vargas, S.A. <i>et al.</i> (1994) c-kit expression by B cell precursors in mouse bone marrow. Stimulation of B cell genesis by in vivo treatment with anti-c-kit antibody. J <u>Immunol. 152 (6): 2845-52.</u>
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign

Informati		Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA1365SBUV605 20471						
Regulato	ry For res							
	d Products	agonte						
	nended Useful Rea EROBLOCK FcR (BUF	-						
MOUSE S	•							
	EROBLOCK FcR (BUF							
MOUSE S	Tel: +1 800 265 7376 Fax: +1 919 878 3751	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50			

Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint