

Datasheet: MCA1365SBY605 BATCH NUMBER 100007457

Description:	RAT ANTI MOUSE CD117:StarBright Yellow 605
Specificity:	CD117
Other names:	C-KIT, SCF-R
Format:	StarBright Yellow 605
Product Type:	Monoclonal Antibody
Clone:	2B8
lsotype:	lgG2b
Quantity:	100 TESTS/0.5ml

Product Details

Applications	derived from testing wi communications from t	thin our laboratorie he originators. Plea al protocol recomm	the following application s, peer-reviewed publica ase refer to references in endations, please visit <u>w</u>	tions or personal ndicated for further
		Yes No	Not Determined	Suggested Dilution
	Flow Cytometry	-		Neat
	necessarily exclude its	use in such proce nmended that the u	or use in a particular tech dures. Suggested workin user titrates the product f e controls.	g dilutions are given as
Target Species	Mouse			
Product Form	Purified IgG conjugate	d to StarBright Yell	ow 605 - liquid	
Max Ex/Em	Fluorophore	Excitation Max (nm) Emission Max (nm)	
	StarBright Yellow 605	572	606	
Preparation	Purified IgG prepared I supernatant	by affinity chromato	graphy on Protein G fro	m tissue culture
Buffer Solution	Phosphate buffered sa	line		
Preservative Stabilisers	0.09% Sodium Azide (l 1% Bovine Serum Albu 0.1% Pluronic F68 0.1% PEG 3350	•		

	0.05% Tween 20
Immunogen	IL-3 dependent mast cells.
External Database Links	UniProt: P05532 Related reagents Entrez Gene: 16590 Kit Related reagents
Synonyms	SI
Fusion Partners	Spleen cells from immunised Wistar rats were fused with cells of the Fox-NY mouse myeloma cell line.
Specificity	Rat anti Mouse CD117 antibody, clone 2B8 recognizes mouse CD117, a ~150 kDa cell surface molecule otherwise known as the Kit, or c-kit, receptor.
	Rat anti Mouse CD117 antibody, clone 2B8 recognizes an epitope within the extracellular region of CD117. In mice, CD117 is predominantly expressed on multipotent haematopoietic stem cells from the adult bone marrow and foetal liver.
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 ⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
Further Reading	 Manova, K. <i>et al.</i> (1992) c-kit receptor and ligand expression in postnatal development of the mouse cerebellum suggests a function for c-kit in inhibitory interneurons. J <u>Neurosci. 12 (12): 4663-76.</u> Maeda, H. <i>et al.</i> (1992) Requirement of c-kit for development of intestinal pacemaker system. <u>Development. 116 (2): 369-75.</u> Takagi, M. <i>et al.</i> (1992) Stimulation of mouse connective tissue-type mast cells by hemopoietic stem cell factor, a ligand for the c-kit receptor. <u>J Immunol. 148 (11): 3446-53.</u> de Vries, P. <i>et al.</i> (1992) Thymus reconstitution by c-kit-expressing hematopoietic stem cells purified from adult mouse bone marrow. <u>J Exp Med. 176 (6): 1503-9.</u> Fleming WH <i>et al.</i> (1993) Steel factor influences the distribution and activity of murine hematopoietic stem cells <i>in vivo</i>. <u>Proc Natl Acad Sci U S A. 90 (8): 3760-4.</u> Rico-Vargas, S.A. <i>et al.</i> (1994) c-kit expression by B cell precursors in mouse bone marrow. Stimulation of B cell genesis by in vivo treatment with anti-c-kit antibody. J <u>Immunol. 152 (6): 2845-52.</u>
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign

Health Ar Information	-		-	sheet documentation #20 Intibodies.com/SDS/MCA	
Regulato	ry	For resea	rch purpose	es only	
	d Product	-	ents		
Recomn	d Products nended Use EROBLOCK FO EROBLOCK FO	ful Reag	<u>1A)</u>		

Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint