

Datasheet: MCA1369BT

Description:	HAMSTER ANTI MOUSE CD11c:Biotin
Specificity:	CD11c
Other names:	INTEGRIN ALPHA X CHAIN
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	N418
Isotype:	IgG
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/5 - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG conjugated to Biotin - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Mouse spleen dendritic cells.

External Database
Links

UniProt:

[Q9QXH4](#) [Related reagents](#)

Entrez Gene:

[16411](#) Itgax [Related reagents](#)

RRID

AB_2129773

Fusion Partners

Spleen cells from immunised Armenian Hamster were fused with cells of the Sp2/0 myeloma cell line.

Specificity

Hamster anti Mouse CD11c antibody, clone N418 recognizes the murine homolog of human CD11c, also known as Integrin Alpha X, a 150/90 kDa member of the beta 2 integrin family. In mice, CD11c is primarily expressed by dendritic cells.

Hamster anti Mouse CD11c antibody, clone N418 has been reported to enhance antigen specific responses when used to target dendritic cells *in vivo* ([Wang et al. 2000](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/B](#)).

References

1. Crowley, M.T. *et al.* (1990) Use of the fluorescence activated cell sorter to enrich dendritic cells from mouse spleen. [J Immunol Methods. 133 \(1\): 55-66.](#)
2. Dahlen, E. *et al.* (1998) Dendritic cells and macrophages are the first and major producers of TNF-alpha in pancreatic islets in the nonobese diabetic mouse. [J Immunol. 160: 3585-93.](#)
3. Wang, H. *et al.* (2000) Rapid antibody responses by low-dose, single-step, dendritic cell-targeted immunization. [Proc Natl Acad Sci U S A. 97 \(2\): 847-52.](#)
4. Lundqvist, J. *et al.* (2010) Concomitant infection decreases the malaria burden but escalates relapsing fever borreliosis. [Infect Immun. 78 \(5\): 1924-30.](#)
5. Beyer, M. *et al.* (2005) The beta2 integrin CD11c distinguishes a subset of cytotoxic pulmonary T cells with potent antiviral effects in vitro and in vivo. [Respir Res. 6: 70.](#)
6. Goupil, M. *et al.* (2009) Macrophage-mediated responses to *Candida albicans* in mice expressing the human immunodeficiency virus type 1 transgene. [Infect Immun. 77: 4136-49.](#)
7. Linehan, S.A.. (2005) The mannose receptor is expressed by subsets of APC in non-lymphoid organs. [BMC Immunol. 6:4.](#)
8. Bjorck, P. (2004) Dendritic cells exposed to herpes simplex virus *in vivo* do not produce IFN-alpha after rechallenge with virus *in vitro* and exhibit decreased T cell alloreactivity. [J Immunol. 172: 5396-404.](#)
9. Mercier, S. *et al.* (2002) Distinct roles of adenovirus vector-transduced dendritic cells, myoblasts, and endothelial cells in mediating an immune response against a transgene product. [J Virol. 76: 2899-911.](#)
10. de Jersey, J. *et al.* (2002) Activation of CD8 T cells by antigen expressed in the pituitary gland. [J Immunol. 169: 6753-9.](#)

11. Dimier-Poisson, I. *et al.* (2003) Protective mucosal Th2 immune response against *Toxoplasma gondii* by murine mesenteric lymph node dendritic cells. [Infect Immun. 71: 5254-65.](#)
12. Gonzalez-Juarrero, M. and Orme, I.M. (2001) Characterization of murine lung dendritic cells infected with *Mycobacterium tuberculosis*. [Infect Immun. 69: 1127-33.](#)
13. Hamada, H. *et al.* (2002) Identification of multiple isolated lymphoid follicles on the antimesenteric wall of the mouse small intestine. [J Immunol. 168: 57-64.](#)
14. Meng, Q. *et al.* (2007) Phenotypes, distribution, and morphological features of antigen-presenting cells in the murine cornea following intravitreal injection. [Mol Vis. 13: 475-86.](#)
15. Moos, M.P. *et al.* (2005) The lamina adventitia is the major site of immune cell accumulation in standard chow-fed apolipoprotein E-deficient mice [Arterioscler Thromb Vasc Biol. 25: 2386-91.](#)
16. Nunez, R. *et al.* (1999) Immortalized cell lines derived from mice lacking both type I and type II IFN receptors unify some functions of immature and mature dendritic cells. [Immunol Cell Biol. 77: 153-63.](#)
17. Ponce, L.V. *et al.* (2005) Adoptive transfer of dendritic cells modulates immunogenesis and tolerogenesis in a neonatal model of murine cutaneous leishmaniasis. [Kinetoplastid Biol Dis. 4: 2.](#)
18. Zhang, L. *et al.* (2011) The inflammatory changes of adipose tissue in late pregnant mice. [J Mol Endocrinol. 47 \(2\): 157-65.](#)
19. Donaldson, D.S. *et al.* (2012) M cell-depletion blocks oral prion disease pathogenesis. [Mucosal Immunol. 5: 216-25.](#)
20. Wada, T. *et al.* (2013) Eplerenone ameliorates the phenotypes of metabolic syndrome with NASH in liver-specific SREBP-1c Tg mice fed high-fat and high-fructose diet. [Am J Physiol Endocrinol Metab. 305 \(11\): E1415-25.](#)
21. Kan, M.J. *et al.* (2015) Arginine deprivation and immune suppression in a mouse model of Alzheimer's disease. [J Neurosci. 35 \(15\): 5969-82.](#)
22. Kayser, B.D. *et al.* (2015) Perinatal Overnutrition Exacerbates Adipose Tissue Inflammation Caused by High-Fat Feeding in C57BL/6J Mice. [PLoS One. 10 \(3\): e0121954.](#)
23. Sehgal, A. *et al.* (2017) c-Rel is dispensable for the differentiation and functional maturation of M cells in the follicle-associated epithelium. [Immunobiology. 222 \(2\): 316-26.](#)
24. Bender, L.H. *et al.* (2020) Intratumoral Administration of a Novel Cytotoxic Formulation with Strong Tissue Dispersive Properties Regresses Tumor Growth and Elicits Systemic Adaptive Immunity in *In Vivo* Models. [Int J Mol Sci. 21 \(12\) Jun 24 \[Epub ahead of print\].](#)
25. Iida, Y. *et al.* (2020) Local injection of CCL19-expressing mesenchymal stem cells augments the therapeutic efficacy of anti-PD-L1 antibody by promoting infiltration of immune cells. [J Immunother Cancer. 8 \(2\) \[Epub ahead of print\].](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M365220:200529'

Printed on 23 Mar 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)