

## Datasheet: MCA1369GA

<b>Description:</b>	HAMSTER ANTI MOUSE CD11c
<b>Specificity:</b>	CD11c
<b>Other names:</b>	INTEGRIN ALPHA X CHAIN
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	N418
<b>Isotype:</b>	IgG
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide

<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Mouse spleen dendritic cells.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">Q9QXH4</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">16411</a> Itgax    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_324695
<b>Fusion Partners</b>	Spleen cells from immunised Armenian Hamster were fused with cells of the Sp2/0 myeloma cell line.
<b>Specificity</b>	<p><b>Hamster anti Mouse CD11c antibody, clone N418</b> recognizes the murine homolog of human CD11c, also known as Integrin Alpha X, a 150/90 kDa member of the beta 2 integrin family. In mice, CD11c is primarily expressed by dendritic cells.</p> <p>Hamster anti Mouse CD11c antibody, clone N418 has been reported to enhance antigen specific responses when used to target dendritic cells <i>in vivo</i> (<a href="#">Wang et al. 2000</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Crowley, M.T. <i>et al.</i> (1990) Use of the fluorescence activated cell sorter to enrich dendritic cells from mouse spleen. <a href="#">J Immunol Methods. 133 (1): 55-66.</a></li> <li>2. Dahlen, E. <i>et al.</i> (1998) Dendritic cells and macrophages are the first and major producers of TNF-alpha in pancreatic islets in the nonobese diabetic mouse. <a href="#">J Immunol. 160: 3585-93.</a></li> <li>3. Wang, H. <i>et al.</i> (2000) Rapid antibody responses by low-dose, single-step, dendritic cell-targeted immunization. <a href="#">Proc Natl Acad Sci U S A. 97 (2): 847-52.</a></li> <li>4. Lundqvist, J. <i>et al.</i> (2010) Concomitant infection decreases the malaria burden but escalates relapsing fever borreliosis. <a href="#">Infect Immun. 78 (5): 1924-30.</a></li> <li>5. Beyer, M. <i>et al.</i> (2005) The beta2 integrin CD11c distinguishes a subset of cytotoxic pulmonary T cells with potent antiviral effects in vitro and in vivo. <a href="#">Respir Res. 6: 70.</a></li> <li>6. Goupil, M. <i>et al.</i> (2009) Macrophage-mediated responses to <i>Candida albicans</i> in mice expressing the human immunodeficiency virus type 1 transgene. <a href="#">Infect Immun. 77: 4136-49.</a></li> <li>7. Linehan, S.A.. (2005) The mannose receptor is expressed by subsets of APC in non-lymphoid organs. <a href="#">BMC Immunol. 6:4.</a></li> <li>8. Bjorck, P. (2004) Dendritic cells exposed to herpes simplex virus <i>in vivo</i> do not produce IFN-alpha after rechallenge with virus <i>in vitro</i> and exhibit decreased T cell alloreactivity. <a href="#">J Immunol. 172: 5396-404.</a></li> <li>9. Mercier, S. <i>et al.</i> (2002) Distinct roles of adenovirus vector-transduced dendritic cells,</li> </ol>

- myoblasts, and endothelial cells in mediating an immune response against a transgene product. [J Virol. 76: 2899-911.](#)
10. de Jersey, J. *et al.* (2002) Activation of CD8 T cells by antigen expressed in the pituitary gland. [J Immunol. 169: 6753-9.](#)
11. Dimier-Poisson, I. *et al.* (2003) Protective mucosal Th2 immune response against *Toxoplasma gondii* by murine mesenteric lymph node dendritic cells. [Infect Immun. 71: 5254-65.](#)
12. Gonzalez-Juarrero, M. and Orme, I.M. (2001) Characterization of murine lung dendritic cells infected with *Mycobacterium tuberculosis*. [Infect Immun. 69: 1127-33.](#)
13. Hamada, H. *et al.* (2002) Identification of multiple isolated lymphoid follicles on the antimesenteric wall of the mouse small intestine. [J Immunol. 168: 57-64.](#)
14. Meng, Q. *et al.* (2007) Phenotypes, distribution, and morphological features of antigen-presenting cells in the murine cornea following intravitreal injection. [Mol Vis. 13: 475-86.](#)
15. Moos, M.P. *et al.* (2005) The lamina adventitia is the major site of immune cell accumulation in standard chow-fed apolipoprotein E-deficient mice [Arterioscler Thromb Vasc Biol. 25: 2386-91.](#)
16. Nunez, R. *et al.* (1999) Immortalized cell lines derived from mice lacking both type I and type II IFN receptors unify some functions of immature and mature dendritic cells. [Immunol Cell Biol. 77: 153-63.](#)
17. Ponce, L.V. *et al.* (2005) Adoptive transfer of dendritic cells modulates immunogenesis and tolerogenesis in a neonatal model of murine cutaneous leishmaniasis. [Kinetoplastid Biol Dis. 4: 2.](#)
18. Zhang, L. *et al.* (2011) The inflammatory changes of adipose tissue in late pregnant mice. [J Mol Endocrinol. 47 \(2\): 157-65.](#)
19. Donaldson, D.S. *et al.* (2012) M cell-depletion blocks oral prion disease pathogenesis. [Mucosal Immunol. 5: 216-25.](#)
20. Wada, T. *et al.* (2013) Eplerenone ameliorates the phenotypes of metabolic syndrome with NASH in liver-specific SREBP-1c Tg mice fed high-fat and high-fructose diet. [Am J Physiol Endocrinol Metab. 305 \(11\): E1415-25.](#)
21. Kan, M.J. *et al.* (2015) Arginine deprivation and immune suppression in a mouse model of Alzheimer's disease. [J Neurosci. 35 \(15\): 5969-82.](#)
22. Kayser, B.D. *et al.* (2015) Perinatal Overnutrition Exacerbates Adipose Tissue Inflammation Caused by High-Fat Feeding in C57BL/6J Mice. [PLoS One. 10 \(3\): e0121954.](#)
23. Sehgal, A. *et al.* (2017) c-Rel is dispensable for the differentiation and functional maturation of M cells in the follicle-associated epithelium. [Immunobiology. 222 \(2\): 316-26.](#)
24. Bender, L.H. *et al.* (2020) Intratumoral Administration of a Novel Cytotoxic Formulation with Strong Tissue Dispersive Properties Regresses Tumor Growth and Elicits Systemic Adaptive Immunity in *In Vivo* Models. [Int J Mol Sci. 21 \(12\) Jun 24 \[Epub ahead of print\].](#)
25. Iida, Y. *et al.* (2020) Local injection of CCL19-expressing mesenchymal stem cells augments the therapeutic efficacy of anti-PD-L1 antibody by promoting infiltration of immune cells. [J Immunother Cancer. 8 \(2\) \[Epub ahead of print\].](#)
26. Kouwenberg, M. *et al.* (2021) Reduced CXCL1 production by endogenous IL-37 expressing dendritic cells does not affect T cell activation. [PLoS One. 16 \(5\): e0251809.](#)
27. Curina, G. *et al.* (2018) Evaluation of immune responses in mice and sheep inoculated with a live attenuated *Brucella melitensis* REV1 vaccine produced in bioreactor. [Vet Immunol Immunopathol. 198: 44-53.](#)

28. Mohanta, S.K. *et al.* (2022) Neuroimmune cardiovascular interfaces control atherosclerosis. [Nature. Apr 27 \[Epub ahead of print\]](#).

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) [DyLight@800](#), [FITC](#)

Goat Anti Hamster IgG (STAR79...) [Biotin](#), [FITC](#), [HRP](#)

### Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL \(MCA2356\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M383041:210513'

**Printed on 23 Jun 2022**

---

© 2022 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)